

A general theory of self-similarity I

Tom Leinster

Department of Mathematics, University of Glasgow
www.maths.gla.ac.uk/~tl
tl@maths.gla.ac.uk

Abstract

Consider a self-similar space X . A typical situation is that X looks like several copies of itself glued to several copies of another space Y , and Y looks like several copies of itself glued to several copies of X , or the same kind of thing with more than two spaces. Thus, the self-similarity of X is described by a system of simultaneous equations. Here I formalize this idea and the notion of a ‘universal solution’ of such a system. I determine exactly when a system has a universal solution and, when one does exist, construct it.

A sequel [Lei2] contains further results and examples, and an introductory article [Lei3] gives an overview.

Contents

Introduction	2
1 Self-similarity systems	5
2 Nondegeneracy	13
3 Coalgebras	20
4 The universal solution	24
5 Set-theoretic proofs	28
6 Topological proofs	38
A Appendix: Solvability	45
References	48

Introduction

Statements of self-similarity are of two kinds: local and global. Local statements say something like ‘almost any small pattern observed in one part of the object can be observed throughout the object, at all scales’. See for instance Chapter 4 of Milnor [Mil], where such statements are made about Julia sets of complex rational functions. Global statements say something like ‘the whole object consists of several smaller copies of itself glued together’; more generally, there may be a whole family of objects, each of which can be described as several objects in the family glued together. Put another way, the contrast is between bottom-up and top-down. This paper and its sequel [Lei2] introduce a theory of global, or top-down, self-similarity.

Statements of self-similarity can also be divided into the glamorous and the mundane. Fractal spaces such as Julia sets, the Cantor set, and Sierpiński’s gasket are certainly self-similar, but there is self-similarity to be found in more everyday objects. For instance, any n -manifold is as locally self-similar as could be: every point is locally isomorphic to every other point. The closed interval $[0, 1]$ is globally self-similar, being isomorphic to two copies of itself glued end to end. We will see that $[0, 1]$ is, in a precise sense, *universal* with this property (Freyd’s Theorem, 1.2). In the same way, barycentric subdivision expresses the topological n -simplex Δ^n as the gluing-together of several smaller copies of itself, and this leads to a universal characterization of the sequence $(\Delta^n)_{n \geq 0}$ among all sequences of topological spaces [Lei2, 2.12].

The shape of this paper can be explained by analogy with linear simultaneous equations.

- a. Consider a sequence $x = (x_1, \dots, x_n)$ of scalars and a system of n simultaneous equations, the i th of which expresses x_i as a linear combination of x_j s (that is, $x_i = \sum_j m_{ij}x_j$).
- b. The coefficients can be encoded as an $n \times n$ matrix M .
- c. The system can be written as $x = Mx$, and we are interested in finding solutions.
- d. We consider only nondegenerate solutions $x \neq 0$.
- e. We may decide to consider only nondegenerate systems of equations, that is, those for which M is nonsingular (x nondegenerate implies Mx nondegenerate).
- f. There are explicit conditions on M equivalent to the existence of a nondegenerate solution of the system (for instance, $\det(M - I) = 0$).
- g. In the case that a nondegenerate solution exists, it can be constructed algorithmically.

Here are the analogous steps for self-similarity equations.

- a. Consider a family $X = (X_a)_{a \in \mathbb{A}}$ of spaces and a system of simultaneous equations, one for each $a \in \mathbb{A}$, expressing each X_a as a gluing-together of X_b s (§1).
- b. The gluing formulas can be encoded as a ‘two-sided $\mathbb{A}$ -module’ M (§1).
- c. The system can be written as $X \cong M \otimes X$. The linear analogy is a simplification: there we had scalars, which are either equal or not, but here we have spaces, which may be equal, isomorphic, or merely connected by a map in one direction or the other. We are interested in ‘ M -coalgebras’, that is, pairs (X, ξ) where $\xi : X \longrightarrow M \otimes X$. We are especially interested in M -coalgebras (X, ξ) possessing a certain universal property. If this property holds then ξ is an isomorphism, so we call such an (X, ξ) a ‘universal solution’ of the system (§§1, 3).
- d. We consider only ‘nondegenerate’ families X (§§1, 2).
- e. We consider only nondegenerate systems of equations, that is, those for which M is a ‘nondegenerate’ module (X nondegenerate implies $M \otimes X$ nondegenerate: §§1, 2).
- f. There is an explicit condition on M equivalent to the existence of a universal solution of the system (§4 and Appendix A; I claim no analogy with determinants).
- g. In the case that a universal solution exists, it can be constructed algorithmically (§§4–6).

The second paper in this series [Lei2] shows how to recognize universal solutions, gives examples, classifies those spaces that are self-similar in at least one way, and uses this classification to reprove some classical results in topology.

Self-similarity is regarded here as *intrinsic* structure: there is no ambient space. (Contrast iterated function systems.) This is like considering abstract groups rather than groups of transformations, or abstract manifolds rather than manifolds embedded in $\mathbb{R}^n$.

I have been referring to ‘spaces’ with self-similar structure. Here and in [Lei2], the types of space considered are sets and general topological spaces. It may be possible to extend the theory to encompass other types of space, hence other types of self-similarity: conformal, statistical, type-theoretic (in the sense of computer science), and so on. The formal mechanism would be to replace the category of sets, which plays a basic role in the theory presented here, by a different category of spaces.

Another long-term goal is to develop the algebraic topology of self-similar spaces, for which the usual homotopical and homological invariants are often useless: in the case of a connected self-similar subset of the plane, for example, they only give us π_1 , which is typically either infinite-dimensional or trivial. However, a description by a set of self-similarity equations is discrete and so might provide useful invariants.

Various other theories are related to this one. Symbolic dynamics [LM] seems most closely related to the case of *discrete* self-similarity systems, studied in §4 of [Lei2]. Iterated function systems ([Fal], [Hut]) are related, but differ crucially in that they take place inside a fixed ambient space; see Examples 2.10 and 2.11 of [Lei2] for more. There is also a paper of Barr [Barr] with obvious similarities to the present work. He discusses terminal coalgebras for an endofunctor, and the metrics associated with them. However, the class of endofunctors he considers is almost disjoint from the class considered here: the categories on which his endofunctors act always have a terminal object, and his terminal coalgebras can be constructed as limits (compare 4.2 below).

Notation and terminology The natural language for this theory is that of categories. The following concepts are used; they are all explained in [Mac].

Let $\mathbb{I}$ be a small category and $\mathcal{E}$ any category. A **diagram of shape $\mathbb{I}$ (or over $\mathbb{I}$)** in $\mathcal{E}$ is a functor $D : \mathbb{I} \longrightarrow \mathcal{E}$. A **cone** on D is an object X of $\mathcal{E}$ together with a map $p_i : X \longrightarrow Di$ for each $i \in \mathbb{I}$, such that $(Du) \circ p_i = p_j$ for any map $u : i \longrightarrow j$ in $\mathbb{I}$. When the cone has a certain universal property, it (or abusively, the object X) is called the **limit** of D . **Cocones** and **colimits** are defined dually.

More generally, let $F : \mathbb{I}^{\text{op}} \times \mathbb{I} \longrightarrow \mathcal{E}$. A **wedge** on F is an object X of $\mathcal{E}$ together with a map $p_i : X \longrightarrow F(i, i)$ for each $i \in \mathbb{I}$, such that $F(1, u) \circ p_i = F(u, 1) \circ p_j$ for any map $u : i \longrightarrow j$ in $\mathbb{I}$. When the wedge has a certain universal property, it (or X) is called the **end** of F , written $X = \int_i F(i, i)$. **Coends** are defined dually and written $X = \int^i F(i, i)$.

(Co)limits are a special case of (co)ends: given $D : \mathbb{I} \longrightarrow \mathcal{E}$, define $F(i, j) = Dj$; then $\int_i F(i, i)$ is the limit of D . So it is reasonable and convenient to write $\int_i Di$ for the limit of D , when it exists. Dually, $\int^i Di$ is the colimit of D .

The sum (coproduct) of a set-indexed family $(D_i)_{i \in I}$ of objects is written $\sum_i D_i$. If $D_i = Y$ for all i then the sum is written $I \times Y$. The sum of a finite family $D_1, \dots, D_n$ of objects is written $D_1 + \dots + D_n$, or 0 if $n = 0$.

Given categories $\mathcal{A}$ and $\mathcal{B}$, the category whose objects are functors from $\mathcal{A}$ to $\mathcal{B}$ and whose morphisms are natural transformations is written $[\mathcal{A}, \mathcal{B}]$.

A **discrete category** is one in which the only maps are the identities. Small discrete categories are therefore just sets.

Top is the category of *all* topological spaces and continuous maps.

The set $\mathbb{N}$ of natural numbers is taken to include 0.

Acknowledgements I am very grateful for the encouragement and help of Clemens Berger, Araceli Bonifant, Edward Crane, Marcelo Fiore, Steve Lack, John Milnor, Justin Sawon, Carlos Simpson, and Ivan Smith. This research has relied crucially on the categories mailing list (see [Fre]); I thank Bob Rosebrugh, who runs it. I am immensely grateful to Jon Nimmo for creating Figure 1-A. The commutative diagrams were made using Paul Taylor's macros.

I gratefully acknowledge a Nuffield Foundation award NUF-NAL 04.




Figure 1-A: (a) The Julia set of $(2z/(1+z^2))^2$; (b), (c) two subsets, rescaled. (Image by Jon Nimmo; see also [Mil, Fig. 2] and [PR, Fig. 53])

1 Self-similarity systems

This section concerns two definitions:

- a ‘self-similarity system’ is a small category $\mathbb{A}$ together with a finite non-degenerate module $M : \mathbb{A} \dashrightarrow \mathbb{A}$
- a ‘universal solution’ of a self-similarity system $(\mathbb{A}, M)$ is a terminal M -coalgebra.

I will explain all this terminology and the underlying ideas using two examples: a Julia set and a theorem of Freyd.

Figure 1-A(a) shows a certain closed subset of the Riemann sphere $\mathbb{C} \cup \{\infty\}$, the Julia set of the function $z \mapsto (2z/(1+z^2))^2$. (What follows is informal, and the reader will not need the definition of Julia set.) Write I_1 for this set, regarded as an abstract topological space. Evidently I_1 has reflectional symmetry in a horizontal axis, so may be written

$$I_1 = \begin{array}{c} I_2 \\ \text{---} \\ I_2 \end{array} \quad (1)$$

where I_2 is a certain space with 4 distinguished points, shown in Figure 1-A(b). In turn, I_2 may be regarded as a gluing-together of subspaces:

$$I_2 = \text{diagram} \quad (2)$$

where I_3 is another space with 4 distinguished points (Figure 1-A(c)). Next, I_3 is two copies of itself glued together:

$$I_3 = \text{diagram} \quad (3)$$

No new spaces appear at this stage, so the process ends. However, the one-point space has played a role (since we are gluing at single points), so I write I_0 for the one-point space and record the trivial equation

$$I_0 = I_0. \quad (4)$$

Equations (1)–(4) are a system of simultaneous equations in which the right-hand sides are ‘two-dimensional formulas’ expressing each object I_n as a gluing of (I_m) s. Informally, a self-similarity system is a system of equations like this, and the particular spaces I_n defined above are a solution of the system.

As a first step towards formalization, observe that the spaces I_n together with their distinguished points define a functor from the category

$$\mathbb{A} = \left(\begin{array}{c} 1 \\ \text{diagram} \\ 0 \end{array} \right)$$

to the category **Set** of sets (or a category of spaces, but let us be conservative for now). This describes the left-hand sides of equations (1)–(4). Next, the gluing formulas define a functor $G : [\mathbb{A}, \mathbf{Set}] \longrightarrow [\mathbb{A}, \mathbf{Set}]$, where if $X \in [\mathbb{A}, \mathbf{Set}]$ then

$$(GX)_1 = \text{diagram} = (X_2 + X_2) / \sim,$$




Figure 1-B: The eight elements of $M(0, 2)$, with three picked out (see (5)).

and similarly for $(GX)_2$, $(GX)_3$, and $(GX)_0$. (The picture of $(GX)_1$ is drawn as if X_0 were a single point.) The simultaneous equations (1)–(4) assert precisely that $I \cong GI$: I is a fixed point of G .

Although these simultaneous equations have many solutions (G has many fixed points), I is in some sense the universal one. This means that the simple diagrams (1)–(4) contain just as much information as the apparently very complex spaces in Figure 1-A: given the system of equations, we recover these spaces as the universal solution. (Caveat: we consider only the intrinsic, topological aspects of self-similar spaces, not how they are embedded into an ambient space or any metric structure.)

Next we have to make rigorous the notion of ‘gluing formula’, so far expressed in pictures. We have a small category $\mathbb{A}$ whose objects index the spaces involved, and I claim that the system of gluing formulas is described by a functor $M : \mathbb{A}^{\text{op}} \times \mathbb{A} \longrightarrow \mathbf{Set}$. The idea is that for $b, a \in \mathbb{A}$,

$$M(b, a) = \{\text{copies of the } b\text{th space used in the gluing formula for the } a\text{th space}\}.$$

For example, in the gluing formula for I_2 , the one-point space I_0 appears 8 times (Figure 1-B), I_1 does not appear at all, I_2 appears twice, and I_3 appears once, so, writing n for an n -element set,

$$M(0, 2) = 8, \quad M(1, 2) = \emptyset, \quad M(2, 2) = 2, \quad M(3, 2) = 1.$$

Similarly, the gluing formula for I_0 is nothing but a single copy of I_0 , so

$$M(0, 0) = 1, \quad M(1, 0) = M(2, 0) = M(3, 0) = \emptyset.$$

For functoriality, consider, for instance, the second of the four maps $0 \longrightarrow 2$. (In the notation above, this determines the second of the four distinguished points of I_2 .) This induces functions

$$M(0, 0) \longrightarrow M(0, 2), \quad M(2, 2) \longrightarrow M(0, 2) \quad (5)$$

(among others). The first function sends the unique element of $M(0, 0)$ to the element of $M(0, 2)$ marked m in Figure 1-B. Writing $M(2, 2) = \{\text{Left}, \text{Right}\}$, the second function sends Left to m' and Right to m'' .

It is convenient to use the language of modules. Given small categories $\mathbb{A}$ and $\mathbb{B}$, a **module** $M : \mathbb{B} \multimap \mathbb{A}$ is a functor $M : \mathbb{B}^{\text{op}} \times \mathbb{A} \longrightarrow \mathbf{Set}$. For example, suppose that $\mathbb{A}$ and $\mathbb{B}$ are monoids (= categories with only one object): then a module $\mathbb{B} \multimap \mathbb{A}$ is a set with a compatible left $\mathbb{A}$ -action and right $\mathbb{B}$ -action. (If we work with categories enriched in abelian groups then one-object categories are rings and modules are bimodules.) Write $b \xrightarrow{m} a$ to mean $m \in M(b, a)$: then a module $M : \mathbb{B} \multimap \mathbb{A}$ is an indexed family $(M(b, a))_{b \in \mathbb{B}, a \in \mathbb{A}}$ of sets together with actions

$$\begin{array}{ccc} b \xrightarrow{m} a & \xrightarrow{f} & a' \\ b' \xrightarrow{g} b & \xrightarrow{m} & a \end{array} \quad \longmapsto \quad \begin{array}{ccc} b & \xrightarrow{fm} & a' \\ b' & \xrightarrow{mg} & a \end{array}$$

such that $(f'f)m = f'(fm)$, $1m = m$, and dually, and $(fm)g = f(mg)$.

(Experts should note that I have reversed the standard convention: a functor $M : \mathbb{B}^{\text{op}} \times \mathbb{A} \longrightarrow \mathbf{Set}$ is written $\mathbb{B} \multimap \mathbb{A}$, not $\mathbb{A} \multimap \mathbb{B}$. One reason is that given such an M , an element $m \in M(b, a)$ can sensibly be written as $b \xrightarrow{m} a$; then with the convention used here, a module $\mathbb{B} \multimap \mathbb{A}$ consists of elements $b \xrightarrow{m} a$.)

The system of self-similarity equations (1)–(4) is therefore encoded by a small category $\mathbb{A}$ and a module $M : \mathbb{A} \multimap \mathbb{A}$. The conceptual distinction between arrows $b \longrightarrow a$ in $\mathbb{A}$ and arrows $b \xrightarrow{m} a$ in M is that the arrows in $\mathbb{A}$ determine where gluing may potentially take place, but the arrows in M say what the gluing actually is.

Here is another way of seeing that the self-similarity equations are encoded by a module. The right-hand side of each of (1)–(4) is a formal gluing of objects of $\mathbb{A}$. ‘Gluing’ are colimits, so if $\widehat{\mathbb{A}}$ is the category obtained by taking $\mathbb{A}$ and freely adjoining all possible colimits then the system of equations amounts to a functor from $\mathbb{A}$ to $\widehat{\mathbb{A}}$. But $\widehat{\mathbb{A}} = [\mathbb{A}^{\text{op}}, \mathbf{Set}]$ (see [MM, I.5.4]), so the system is a functor $\mathbb{A} \longrightarrow [\mathbb{A}^{\text{op}}, \mathbf{Set}]$, that is, a module $\mathbb{A} \multimap \mathbb{A}$.

Simple colimits can often be described by diagrams or formulas, which provides a useful way of specifying simple self-similarity systems. For instance, equations (1)–(4) are an informal description of our $(\mathbb{A}, M)$. In the same way, the self-similarity system $(\mathbb{A}, M)$ in which $\mathbb{A}$ is the discrete category with object-set $\{0, 1\}$ and $M(0, 0) = M(0, 1) = M(1, 1) = 1$ and $M(1, 0) = \emptyset$ can informally (and more intelligibly) be described by the equations

$$\begin{array}{lcl} A & = & A \\ B & = & A + B. \end{array} \tag{6}$$

Given rings A, B and C , an (A, B) -bimodule M , and a (B, C) -bimodule N , there arises an (A, C) -bimodule $M \otimes_B N$. There is a similar tensor product of categorical modules:

$$C \xrightarrow{N} B \xrightarrow{M} A \quad \text{gives rise to} \quad C \xrightarrow{M \otimes N} A.$$

Here $M \otimes N$ is defined by the coend formula

$$(M \otimes N)(c, a) = \int^b M(b, a) \times N(c, b).$$

(See [Mac, Ch. IX] for an explanation of coends.) Concretely,

$$(M \otimes N)(c, a) = \left(\sum_b M(b, a) \times N(c, b) \right) / \sim$$

where $\sim$ is the equivalence relation generated by $(mg, n) \sim (m, gn)$ for all m, g, n of the appropriate types; the element of $(M \otimes N)(c, a)$ represented by $(m, n) \in M(b, a) \times N(c, b)$ is written $m \otimes n$. The tensor product of modules is associative and unital up to coherent isomorphism. (More precisely, categories, modules, and their maps form a bicategory: [Bor, 7.8.2].)

In the Julia set example, the endofunctor G of $[\mathbb{A}, \mathbf{Set}]$ is just $M \otimes -$. This makes sense: a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ can be regarded as a module $X : \mathbf{1} \dashrightarrow \mathbb{A}$ (where $\mathbf{1}$ is the category with one object and only the identity arrow), and there is then a tensor product $M \otimes X : \mathbf{1} \dashrightarrow \mathbb{A}$, that is, a functor $M \otimes X : \mathbb{A} \longrightarrow \mathbf{Set}$. It is given by

$$(M \otimes X)a = \int^b M(b, a) \times Xb = \left(\sum_b M(b, a) \times Xb \right) / \sim. \quad (7)$$

In the example, taking $a = 2$, this says that

$$(M \otimes X)_2 = (8 \times X_0 + 2 \times X_2 + X_3) / \sim$$

where $\sim$ identifies the various copies of X_0 with their images in X_2 and X_3 . In general, (7) makes precise the idea that $M(b, a)$ is the set of copies of the b th space used in the gluing formula for the a th space.


The second example is a result of Peter Freyd, and comes from a very different direction. To state it we need some more terminology.

Given a category $\mathcal{C}$ and an endofunctor G of $\mathcal{C}$, a **G -coalgebra** is an object X of $\mathcal{C}$ together with a map $\xi : X \longrightarrow GX$. (For instance, if $\mathcal{C}$ is a category of modules and $GX = X \otimes X$ then a G -coalgebra is a coalgebra—not necessarily coassociative—in the usual sense.) A **map** $(X, \xi) \longrightarrow (X', \xi')$ of coalgebras is a map $X \longrightarrow X'$ in $\mathcal{C}$ making the evident square commute. Depending on what G is, the category of G -coalgebras may or may not have a terminal object, but if it does then it is a fixed point:


Lemma 1.1 (Lambek [Lam]) *Let $\mathcal{C}$ be a category and G an endofunctor of $\mathcal{C}$. If (I, ι) is terminal in the category of G -coalgebras then $\iota : I \longrightarrow GI$ is an isomorphism.* $\square$

Here is what Freyd said, strengthened slightly. Let $\mathcal{C}$ be the category whose objects are diagrams $X_0 \xrightleftharpoons[v]{u} X_1$ where X_0 and X_1 are sets and u and v are


injections with disjoint images; then an object of $\mathcal{C}$ can be drawn as



where the copies of X_0 on the left and the right are the images of u and v respectively. A map $X \longrightarrow X'$ in $\mathcal{C}$ consists of functions $X_0 \longrightarrow X'_0$ and $X_1 \longrightarrow X'_1$ making the evident two squares commute. Now, given $X \in \mathcal{C}$ we can form a new object GX of $\mathcal{C}$ by gluing two copies of X end to end:



Formally, the endofunctor G of $\mathcal{C}$ is defined by pushout:



For example, the unit interval with its endpoints distinguished forms an object

$$I = \left(\{\star\} \xrightleftharpoons[1]{0} [0, 1] \right)$$

of $\mathcal{C}$, and GI is naturally described as an interval of length 2, again with its endpoints distinguished:

$$GI = \left(\{\star\} \xrightleftharpoons[2]{0} [0, 2] \right).$$

So there is a coalgebra structure $\iota : I \longrightarrow GI$ on I given by multiplication by two. Freyd's Theorem says that this is, in fact, the *universal* example of a G -coalgebra:

Theorem 1.2 (Freyd [Fre]) (I, ι) is terminal in the category of G -coalgebras.

This follows from a general result [Lei2, 2.1]. A direct proof runs roughly as follows. Take a G -coalgebra (X, ξ) and an element $x_0 \in X_1$. Then $\xi(x_0) \in (GX)_1$ is in either the left-hand or the right-hand copy of X_1 , so gives rise to a binary digit $m_1 \in \{0, 1\}$ and a new element $x_1 \in X_1$. (If $\xi(x_0)$ is in the overlap between the two copies of X_1 , choose left or right arbitrarily.) Iterating gives a binary representation $0.m_1m_2\dots$ of an element of $[0, 1]$, and this is the image of x_0 under the unique coalgebra map $(X, \xi) \longrightarrow (I, \iota)$.

The formalism introduced in the Julia set example can also be used here. The category $\mathcal{C}$ is a full subcategory of $[\mathbb{A}, \mathbf{Set}]$, where $\mathbb{A} = (0 \xrightleftharpoons[\tau]{\sigma} 1)$. The module $M : \mathbb{A} \multimap \mathbb{A}$ corresponding to the gluing formula (8) is given by

$$\begin{array}{ccc} M(-, 0) & \xrightleftharpoons[\tau \cdot -]{\sigma \cdot -} & M(-, 1) \\ \\ \begin{array}{ccc} M(0, -) & \begin{array}{c} \{id\} \xrightleftharpoons[1]{0} \{0, \frac{1}{2}, 1\} \\ \uparrow \uparrow \\ \emptyset \xrightarrow{\quad} \{[0, \frac{1}{2}], [\frac{1}{2}, 1]\} \end{array} \\ \uparrow \uparrow \quad \uparrow \uparrow \\ \begin{array}{ccc} \vdash \cdot \sigma & & \vdash \cdot \tau \end{array} \end{array} & & \begin{array}{ccc} \inf & & \sup \end{array} \end{array} \quad (9)$$

(Here $M(0, 1)$ is just a 3-element set and $M(1, 1)$ a 2-element set, but their elements have been named suggestively.) Then $M \otimes -$ defines an endofunctor of $[\mathbb{A}, \mathbf{Set}]$, and the endofunctor G of $\mathcal{C}$ is its restriction.

The only remaining mystery is the condition that the functions $u, v : X_0 \longrightarrow X_1$ are injections with disjoint images. Without it, the theorem would degenerate entirely, as the terminal coalgebra would be $(\{\star\} \rightrightarrows \{\star\})$. It turns out to be a form of flatness.

First note that any two functors $X : \mathbb{A} \longrightarrow \mathbf{Set}$ and $Y : \mathbb{A}^{op} \longrightarrow \mathbf{Set}$ on a small category $\mathbb{A}$ have a tensor product $Y \otimes X$ (a mere set), since they can be regarded as modules

$$\mathbf{1} \multimap \mathbb{A} \xrightarrow{X} \mathbb{A} \xrightarrow{Y} \mathbf{1}.$$

Explicitly,

$$Y \otimes X = \int^a Y a \times X a = \left(\sum_a Y a \times X a \right) / \sim.$$

By definition, a left module X over a ring is flat if the functor $- \otimes X$ preserves finite limits. There is an analogous definition when X is a $\mathbf{Set}$ -valued functor, but we actually want something weaker:

Definition 1.3 Let $\mathbb{A}$ be a small category. A functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is **componentwise flat** or **nondegenerate** if the functor

$$- \otimes X : [\mathbb{A}^{op}, \mathbf{Set}] \longrightarrow \mathbf{Set}$$

preserves finite connected limits. The full subcategory of $[\mathbb{A}, \mathbf{Set}]$ consisting of the nondegenerate functors is written $[\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$.

(A category is **connected** if it is nonempty and cannot be written as a coproduct of two nonempty categories; a **finite connected limit** is a limit over a finite connected category.)

This definition is explained at length in §2. It is shown there that a $\mathbf{Set}$ -valued functor X on $(0 \xrightleftharpoons[\tau]{\sigma} 1)$ is nondegenerate exactly when the functions

$X\sigma$ and $X\tau$ are injections with disjoint images: so $\mathcal{C} = [\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$. Proposition 3.1 says that for general $\mathbb{A}$ and M , the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Set}]$ restricts to an endofunctor of $[\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$ as long as M is nondegenerate:

Definition 1.4 Let $\mathbb{A}$ and $\mathbb{B}$ be small categories. A module $M : \mathbb{B} \multimap \mathbb{A}$ is **nondegenerate** if $M(b, -) : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate for each $b \in \mathbb{B}$.

It is visible from diagram (9) that the Freyd module M is nondegenerate.

To summarize: starting with a certain small category $\mathbb{A}$ and a certain nondegenerate module $M : \mathbb{A} \multimap \mathbb{A}$, Freyd's Theorem describes the terminal coalgebra for the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$. The Julia set example uses a different $\mathbb{A}$ and M . In general, I will restrict to those $\mathbb{A}$ and M for which 'each gluing formula is finite', although I allow there to be infinitely many such formulas (infinitely many objects of $\mathbb{A}$).

To make this precise, recall that any presheaf $Y : \mathbb{A}^{\text{op}} \longrightarrow \mathbf{Set}$ on a small category $\mathbb{A}$ has a **category of elements** $\mathbb{E}(Y)$, whose objects are pairs (a, y) with $a \in \mathbb{A}$ and $y \in Ya$ and whose maps $(a, y) \longrightarrow (a', y')$ are maps $f : a \longrightarrow a'$ in $\mathbb{A}$ such that $(Yf)y' = y$. Similarly, any covariant functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ has a **category of elements** $\mathbb{E}(X)$. In each case, there is a covariant projection functor from the category of elements to $\mathbb{A}$.

A module $M : \mathbb{A} \multimap \mathbb{A}$ is **finite** if for each $a \in \mathbb{A}$, the category $\mathbb{E}(M(-, a))$ is finite. Explicitly, this says that for each $a \in \mathbb{A}$ there are only finitely many diagrams of the form

$$b' \xrightarrow{f} b \multimap^m a.$$

Certainly this holds if, as in the Freyd example, the category $\mathbb{A}$ and the sets $M(b, a)$ are finite.

Definition 1.5 A **self-similarity system** is a small category $\mathbb{A}$ together with a finite nondegenerate module $M : \mathbb{A} \multimap \mathbb{A}$.

Definition 1.6 Let $(\mathbb{A}, M)$ be a self-similarity system. An **M -coalgebra (in $\mathbf{Set}$)** is a coalgebra for the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$. A **universal solution** of $(\mathbb{A}, M)$ (in $\mathbf{Set}$) is a terminal object in the category of M -coalgebras.

In this language, Freyd's Theorem describes the universal solution of a certain self-similarity system.

There is also a topological version of Freyd's Theorem. One's first thought might be to take the definitions of $\mathcal{C}$ and G and change 'set' to 'space' and 'function' to 'continuous map'; but then the universal solution is given by the indiscrete topology on $[0, 1]$. The Euclidean topology appears, however, if we insist that $u, v : X_0 \longrightarrow X_1$ are closed maps. So, let $\mathcal{C}'$ be the category whose objects are diagrams $X_0 \rightrightarrows X_1$ of topological spaces and continuous closed injections with disjoint images, define an endofunctor G' of $\mathcal{C}'$ just as G was defined, and define (I, ι) as before, with the Euclidean topology on $[0, 1]$. Then:

Theorem 1.7 (Topological Freyd) (I, ι) is terminal in the category of G' -coalgebras.

(See [Lei2, 2.1] for a proof.) This suggests the following general definition.

Definition 1.8 Let $\mathbb{A}$ be a small category. Write $U : \mathbf{Top} \longrightarrow \mathbf{Set}$ for the underlying set functor. A functor $X : \mathbb{A} \longrightarrow \mathbf{Top}$ is **nondegenerate** if $U \circ X$ is nondegenerate and for each map f in $\mathbb{A}$, the map Xf is closed. The full subcategory of $[\mathbb{A}, \mathbf{Top}]$ formed by the nondegenerate functors is written $[\mathbb{A}, \mathbf{Top}]_{\text{nondegen}}$.

We will often meet functors X for which each space Xa is compact Hausdorff, and then the closedness condition is automatically satisfied.

We also want a general notion of topological M -coalgebra. Let $\mathcal{E}$ be a category with finite colimits, $\mathbb{A}$ a small category, and $M : \mathbb{A} \dashrightarrow \mathbb{A}$ a finite module. Then there is an endofunctor $M \otimes -$ of $[\mathbb{A}, \mathcal{E}]$ defined by the usual coend formula

$$(M \otimes X)a = \int^b M(b, a) \times Xb.$$

Proposition 3.5 says that for any self-similarity system $(\mathbb{A}, M)$, the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Top}]$ restricts to an endofunctor of $[\mathbb{A}, \mathbf{Top}]_{\text{nondegen}}$.

Definition 1.9 Let $(\mathbb{A}, M)$ be a self-similarity system. An M -**coalgebra in $\mathbf{Top}$** is a coalgebra for the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Top}]_{\text{nondegen}}$. A **universal solution of $(\mathbb{A}, M)$ in $\mathbf{Top}$** is a terminal M -coalgebra in $\mathbf{Top}$.

For example, the topological Freyd theorem describes the universal solution in $\mathbf{Top}$ of a certain self-similarity system.

Universal solutions are evidently unique (up to canonical isomorphism) when they exist. The word ‘solution’ is justified by Lambek’s Lemma (1.1): if (J, γ) is a universal solution then $M \otimes J \cong J$. Note that the converse fails: the empty functor $J = \emptyset$ has a unique coalgebra structure, is nondegenerate, and satisfies $M \otimes J \cong J$, but is not usually the terminal coalgebra. (It is the initial algebra.) In the Freyd interval example, there are many coalgebras (J, γ) such that γ is an isomorphism but (J, γ) is not the universal solution: for instance, the universal solution can be multiplied by any space S to give such a coalgebra (J, γ) , with $J = (S \rightrightarrows [0, 1] \times S)$.

Just as an ordinary system of equations need not have a solution, a self-similarity system need not have a universal solution. In §4 we meet an explicit condition equivalent to the existence of a universal solution.

2 Nondegeneracy

In this section I explain nondegeneracy, first by theory and then by examples. The theory leads up to the result that a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate if and only if it satisfies the following explicit conditions:

ND1 given

$$\begin{array}{ccc} a & & a' \\ & \searrow f & \swarrow f' \\ & b & \end{array}$$

in $\mathbb{A}$ and $x \in Xa$, $x' \in Xa'$ such that $fx = f'x'$, there exist a commutative square

$$\begin{array}{ccc} & c & \\ g \swarrow & & \searrow g' \\ a & & a' \\ f \searrow & & \swarrow f' \\ & b & \end{array}$$

and $z \in Xc$ such that $x = gz$, $x' = g'z$

ND2 given $a \xrightleftharpoons[f']{f} b$ in $\mathbb{A}$ and $x \in Xa$ such that $fx = f'x$, there exist a fork

$$c \xrightarrow{g} a \xrightleftharpoons[f']{f} b \quad (10)$$

and $z \in Xc$ such that $x = gz$. (A diagram (10) is a **fork** if $fg = f'g$.)

The examples illustrate that nondegeneracy means ‘no unforced equalities’.

For the theory I will assume some more sophisticated categorical knowledge than in the rest of the paper. Readers who prefer to take it on trust can jump to the passage after Corollary 2.9.

None of this theory is new: it goes back to Grothendieck and Verdier [GV] and Gabriel and Ulmer [GU], and was later developed by Weberpals [Web], Lair [Lai], Ageron [Age], and Adámek, Borceux, Lack, and Rosický [ABLR]. More general statements of much of what follows can be found in [ABLR].

Let us begin with ordinary flat functors. A functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ on a small category $\mathbb{A}$ is **flat** if $- \otimes X : [\mathbb{A}^{\text{op}}, \mathbf{Set}] \longrightarrow \mathbf{Set}$ preserves finite limits.

Theorem 2.1 (Flatness) *Let $\mathbb{A}$ be a small category. The following conditions on a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ are equivalent:*

- a. *X is flat*
- b. *every finite diagram in $\mathbb{E}(X)$ admits a cone*
- c. *each of the following holds:*
 - *there exists $a \in \mathbb{A}$ for which Xa is nonempty*
 - *given $a, a' \in \mathbb{A}$, $x \in Xa$, and $x' \in Xa'$, there exist a diagram*

$$a \xleftarrow{g} c \xrightarrow{g'} a' \text{ in } \mathbb{A} \text{ and } z \in Xc \text{ such that } gz = x \text{ and } g'z = x'$$
 - **ND2.**

Proof See [Bor, §6.3] or [MM, VII.6], for instance. $\square$

The following lemmas are often used to prove this theorem and will also be needed later.

Lemma 2.2 (Existence of cones) *Let $\mathbb{I}$ and $\mathbb{A}$ be small categories and let $X : \mathbb{A} \longrightarrow \mathbf{Set}$. If $- \otimes X : [\mathbb{A}^{\text{op}}, \mathbf{Set}] \longrightarrow \mathbf{Set}$ preserves limits of shape $\mathbb{I}$ then every diagram of shape $\mathbb{I}$ in $\mathbb{E}(X)$ admits a cone.*

Remark The hypothesis can be weakened to say that $- \otimes X$ preserves limits of shape $\mathbb{I}$ of *diagrams of representables*, that is, of diagrams $\mathbb{I} \longrightarrow [\mathbb{A}^{\text{op}}, \mathbf{Set}]$ that factor through the Yoneda embedding of $\mathbb{A}$.

Proof Let $D : \mathbb{I} \longrightarrow \mathbb{E}(X)$ be a diagram of shape $\mathbb{I}$, writing $Di = (a_i, x_i)$ for each $i \in \mathbb{I}$. Then there is a diagram $Y_\bullet : \mathbb{I} \longrightarrow [\mathbb{A}^{\text{op}}, \mathbf{Set}]$ given by $Y_i = \mathbb{A}(-, a_i)$, so by hypothesis the canonical map

$$\int^a \left(\int_i \mathbb{A}(a, a_i) \right) \times Xa \cong \left(\int_i Y_i \right) \otimes X \longrightarrow \int_i (Y_i \otimes X) \cong \int_i Xa_i$$

is a bijection, and in particular a surjection. Since $(x_i)_{i \in \mathbb{I}} \in \int_i Xa_i$, there exist $a \in \mathbb{A}$ and

$$((p_i)_{i \in \mathbb{I}}, x) \in \left(\int_i \mathbb{A}(a, a_i) \right) \times Xa$$

such that $p_i x = x_i$ for all i . Hence $\left((a, x) \xrightarrow{p_i} (a_i, x_i) \right)_{i \in \mathbb{I}}$ is a cone on D . $\square$

Say that a category $\mathcal{C}$ has the **square-completion property** if there exists a cone on every diagram of shape $(\bullet \longrightarrow \bullet \longleftarrow \bullet)$ in $\mathcal{C}$.

Lemma 2.3 (Connectedness by spans) *Two objects c, c' of a category with the square-completion property are in the same connected-component if and only if there exists a span $c \longleftarrow c'' \longrightarrow c'$ connecting them.* $\square$

Lemma 2.4 (Equality in a tensor product) *Let $\mathbb{A}$ be a small category and*

$$X : \mathbb{A} \longrightarrow \mathbf{Set}, \quad Y : \mathbb{A}^{\text{op}} \longrightarrow \mathbf{Set}.$$

Suppose that $\mathbb{E}(X)$ has the square-completion property. Let

$$a, a' \in \mathbb{A}, \quad (y, x) \in Ya \times Xa, \quad (y', x') \in Ya' \times Xa'.$$

Then $y \otimes x = y' \otimes x'$ if and only if there exists a span $a \xleftarrow{f} b \xrightarrow{f'} a'$ and an element $z \in Xb$ such that $x = fz$, $x' = f'z$, and $yf = y'f'$.

Proof See the remarks after the statement of Theorem VII.6.3 in [MM]. $\square$

There is a characterization of componentwise flat functors very similar to that of flat functors in Theorem 2.1. First we need a fact about connectedness.

Lemma 2.5 (Components of a functor) *Any functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ on a small category $\mathbb{A}$ can be written as a sum $X \cong \sum_{j \in J} X_j$ where J is some set and $\mathbb{E}(X_j)$ is connected for each $j \in J$.*

Proof We use the equivalence between **Set**-valued functors and discrete opfibrations. Write $\mathbb{E}(X)$ as a sum $\sum_{j \in J} \mathbb{E}_j$ of connected categories. For each j , the restriction to $\mathbb{E}_j$ of the projection $\mathbb{E}(X) \longrightarrow \mathbb{A}$ is still a discrete opfibration, so corresponds to a functor $X_j : \mathbb{A} \longrightarrow \mathbf{Set}$. Then

$$\mathbb{E} \left(\sum X_j \right) \cong \sum \mathbb{E}(X_j) \cong \sum \mathbb{E}_j \cong \mathbb{E}(X)$$

compatibly with the projections, so $\sum X_j \cong X$. $\square$

Here is the main result.

Theorem 2.6 (Componentwise flatness) *Let $\mathbb{A}$ be a small category. The following conditions on a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ are equivalent:*

- a. X is componentwise flat
- b. every finite connected diagram in $\mathbb{E}(X)$ admits a cone
- c. X satisfies **ND1** and **ND2**
- d. X is a sum of flat functors.

Remark In Lemma 2.5, the functors X_j may be regarded as the connected-components of X . A further equivalent condition is that every connected-component of X is flat: hence the name ‘componentwise flat’.

Proof

(a) $\Rightarrow$ (b) Follows from Lemma 2.2.

(b) $\Rightarrow$ (c) **ND1** says that every diagram of shape $(\bullet \longrightarrow \bullet \longleftarrow \bullet)$ in $\mathbb{E}(X)$ admits a cone, and similarly **ND2** for $(\bullet \rightrightarrows \bullet)$.

(c) $\Rightarrow$ (d) Write $X \cong \sum_{j \in J} X_j$ as in Lemma 2.5. Then in each $\mathbb{E}(X_j)$, there exists a cone on every diagram of shape

$$(\bullet \longrightarrow \bullet \longleftarrow \bullet) \quad \text{or} \quad (\bullet \rightrightarrows \bullet)$$

(since $\mathbb{E}(X) \cong \sum_j \mathbb{E}(X_j)$), of shape $\emptyset$ (since $\mathbb{E}(X_j)$ is connected and therefore nonempty), and of shape $(\bullet \quad \bullet)$ (since $\mathbb{E}(X_j)$ is connected and has the square-completion property). So by (c) $\Rightarrow$ (a) of Theorem 2.1, each X_j is flat.

(d) $\Rightarrow$ (a) Any sum of componentwise flat functors is componentwise flat, as follows from the fact that sums commute with connected limits in **Set**. $\square$

Corollary 2.7 (Componentwise filtered categories) *The following conditions on a small category $\mathbb{B}$ are equivalent:*

- a. finite connected limits commute with colimits of shape $\mathbb{B}$ in **Set**
- b. every finite connected diagram in $\mathbb{B}$ admits a cocone
- c. every diagram $b_1 \longleftarrow b_3 \longrightarrow b_2$ in $\mathbb{B}$ can be completed to a commutative square, and every parallel pair $b_1 \xrightarrow{f} b_2$ of arrows in $\mathbb{B}$ can be extended to a cofork.

Proof In Theorem 2.6, take $\mathbb{A} = \mathbb{B}^{\text{op}}$ and X to be the functor with constant value 1. Then $\mathbb{E}(X) \cong \mathbb{B}^{\text{op}}$ and $- \otimes X$ forms colimits. The result follows. $\square$

A small category $\mathbb{B}$ satisfying the equivalent conditions of Corollary 2.7 is called **componentwise filtered**. (Grothendieck and Verdier say ‘pseudo-filtrante’ [GV], and a further equivalent condition is that each connected-component is filtered.) So $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is componentwise flat just when $\mathbb{E}(X)$ is componentwise cofiltered.

Componentwise flatness relates to limit preservation as follows.

Lemma 2.8 *Componentwise flat functors preserve finite connected limits.*

Proof Let $\mathbb{A}$ be a small category and $X : \mathbb{A} \longrightarrow \mathbf{Set}$ a componentwise flat functor. We have

$$X \cong \left(\mathbb{A} \xrightarrow{\text{Yoneda}} [\mathbb{A}^{\text{op}}, \mathbf{Set}] \xrightarrow{- \otimes X} \mathbf{Set} \right) \quad (11)$$

and the Yoneda embedding preserves limits. $\square$

Corollary 2.9 *Let $\mathbb{A}$ be a small category with all pullbacks and equalizers. Then a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is componentwise flat if and only if it preserves pullbacks and equalizers.*

Proof Suppose that X preserves pullbacks and equalizers. By (11), $- \otimes X$ preserves pullbacks and equalizers of representables, so by the Remark after the statement of Lemma 2.2, any diagram of shape $(\bullet \longrightarrow \bullet \longleftarrow \bullet)$ or $(\bullet \rightrightarrows \bullet)$ in $\mathbb{E}(X)$ admits a cone. But this says that X satisfies **ND1** and **ND2**. $\square$

Intuitively, nondegeneracy (componentwise flatness) of a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ says that no equation between elements of X holds unless it must. For example, if $\mathbb{A} = (\bullet \longrightarrow \bullet)$ then a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is a map $(X_0 \xrightarrow{i} X_1)$ of sets, and nondegeneracy of X says that the equation $ix_0 = ix'_0$ holds only when it must, that is, only when $x_0 = x'_0$; thus, X is nondegenerate just when i is injective (Example 2.10). Or, let $\mathbb{A}$ be the category generated by objects and arrows

$$0 \xrightleftharpoons[\tau]{\sigma} 1 \xrightarrow{\rho} 2 \quad (12)$$

subject to $\rho\sigma = \rho\tau$. If $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate then the equation $(X\rho)x_1 = (X\rho)x'_1$ holds only when it must, that is, when $x_1 = x'_1$ or there exists x_0 satisfying $\{x_1, x'_1\} = \{(X\sigma)x_0, (X\tau)x_0\}$ (Example 2.12).

Let us work out what nondegeneracy says for various specific categories $\mathbb{A}$. Note that **ND1** holds automatically if either f or f' is an isomorphism, and that **ND2** holds automatically if $f = f'$; we therefore ignore these cases. Moreover, if f is monic then **ND1** in the case $f = f'$ just says that Xf is injective; indeed, we already know from Lemma 2.8 that nondegenerate functors preserve monics.

Example 2.10 Let $\mathbb{A} = (0 \xrightarrow{\sigma} 1)$. Then $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate if and only if the function $X\sigma : X0 \longrightarrow X1$ is injective.

Example 2.11 Let $\mathbb{A} = \left(0 \begin{smallmatrix} \xrightarrow{\sigma} \\ \xrightarrow{\tau} \end{smallmatrix} 1\right)$, so that a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is a pair $\left(X_0 \begin{smallmatrix} \xrightarrow{X\sigma} \\ \xrightarrow{X\tau} \end{smallmatrix} X_1\right)$ of functions. Then **ND1** in the case $f = f'$ says that $X\sigma$ and $X\tau$ are injective. The only other nontrivial case of **ND1** is $f = \sigma, f' = \tau$, and since the diagram

$$\begin{array}{ccc} 0 & & 0 \\ & \searrow \sigma & \swarrow \tau \\ & 1 & \end{array}$$

cannot be completed to a commutative square, **ND1** says that $X\sigma$ and $X\tau$ have disjoint images. The only nontrivial case of **ND2** is $f = \sigma, f' = \tau$, and since the diagram $\left(0 \begin{smallmatrix} \xrightarrow{\sigma} \\ \xrightarrow{\tau} \end{smallmatrix} 1\right)$ cannot be completed to a fork, this says that $(X\sigma)x_0 \neq (X\tau)x_0$ for all $x_0 \in X_0$, which we already know. So a nondegenerate functor on $\mathbb{A}$ is a parallel pair of injections with disjoint images, as claimed in §1.

Example 2.12 Let $\mathbb{A}$ be the category generated by objects and arrows (12) subject to $\rho\sigma = \rho\tau$, and consider a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$. The nontrivial cases of **ND1** are:

- $f, f' \in \{\sigma, \tau\}$: then as in Example 2.11, **ND1** says that $X\sigma$ and $X\tau$ are injections with disjoint images
- $f = f' = \rho$: the pairs of maps completing the diagram

$$\begin{array}{ccc} 1 & & 1 \\ & \searrow \rho & \swarrow \rho \\ & 2 & \end{array}$$

to a commutative square are

$$(\text{id}, \text{id}), (\sigma, \sigma), (\tau, \tau), (\sigma, \tau), (\tau, \sigma),$$

so **ND1** says that if $x_1, x'_1 \in X_1$ and $(X\rho)x_1 = (X\rho)x'_1$ then $x_1 = x'_1$ (first three cases) or there exists $x_0 \in X_0$ such that $x_1 = (X\sigma)x_0$ and $x'_1 = (X\tau)x_0$ (fourth case) or *vice versa* (fifth case).

- $f = f' = \rho\sigma$: then **ND1** says that $X(\rho\sigma)$ is injective
- $f = \rho, f' = \rho\sigma$: this can be seen to be redundant.

The only nontrivial case of **ND2** is $f = \sigma, f' = \tau$, and as we saw in Example 2.11, this too is redundant. So X is nondegenerate just when:

- $X\sigma, X\tau$, and $X(\rho\sigma)$ are injective
- $X\sigma$ and $X\tau$ have disjoint images

- if $(X\rho)x_1 = (X\rho)x'_1$ then $x_1 = x'_1$ or there exists x_0 such that $\{x_1, x'_1\} = \{(X\sigma)x_0, (X\tau)x_0\}$.

An example of a nondegenerate functor on $\mathbb{A}$ is the diagram

$$\{\star\} \begin{array}{c} \xrightarrow{0} \\ \xrightarrow{1} \end{array} [0, 1] \longrightarrow S^1$$

exhibiting the circle as an interval with its endpoints identified.

Example 2.13 Let $\mathbb{A}$ be the category generated by objects and arrows

$$0 \begin{array}{c} \xrightarrow{\sigma_1} \\ \xrightarrow{\tau_1} \end{array} 1 \begin{array}{c} \xrightarrow{\sigma_2} \\ \xrightarrow{\tau_2} \end{array} \dots$$

subject to $\sigma_{k+1}\sigma_k = \tau_{k+1}\sigma_k$ and $\sigma_{k+1}\tau_k = \tau_{k+1}\tau_k$ for all $k \geq 1$. A functor $\mathbb{A}^{\text{op}} \longrightarrow \mathbf{Set}$ is usually called a globular set or an ω -graph. It can be shown that a coglobular set $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate precisely when

- for all $k \geq 1$, $X\sigma_k$ and $X\tau_k$ are injective
- for all $k \geq 1$ and $x, x' \in X_k$ satisfying $(X\sigma_{k+1})x = (X\tau_{k+1})x'$, we have $x = x' \in \text{image}(X\sigma_k) \cup \text{image}(X\tau_k)$
- the images of $X\sigma_1$ and $X\tau_1$ are disjoint.

For instance, the underlying coglobular set of any disk in the sense of Joyal ([Joy], [Lei1]) is nondegenerate.

We finish this section with a diagrammatic formulation of nondegeneracy of a module. This will be invaluable later.

First observe that the notion of commutative diagram in a category $\mathbb{A}$ can be extended to include elements of a module $M : \mathbb{A} \dashrightarrow \mathbb{A}$. For instance, the diagram

$$\begin{array}{ccccc} a_2 & \xrightarrow{m_2} & a_1 & \xrightarrow{m_1} & a_0 \\ f_2 \downarrow & & \downarrow f_1 & & \downarrow f_0 \\ a'_2 & \xrightarrow{m'_2} & a'_1 & \xrightarrow{m'_1} & a'_0 \end{array}$$

is said to **commute** if $m'_2 f_2 = f_1 m_2$ and $m'_1 f_1 = f_0 m_1$. (We never attempt to compose paths containing more than one crossed arrow $\dashrightarrow$.) Similarly, the diagram

$$\begin{array}{ccc} & \xrightarrow{m} & \\ a_1 & \dashrightarrow & a_0 \\ f_1 \downarrow & g_1 \downarrow & f_0 \downarrow \\ & \dashrightarrow & \\ a'_1 & \dashrightarrow & a'_0 \\ & m' & \end{array}$$


commutes serially if $m'f_1 = f_0m$ and $m'g_1 = g_0m$, and the diagram

$$b \xrightarrow{m} a \xrightleftharpoons[f']{f} c$$

is a **fork** if $fm = f'm$.


In this language, $M : \mathbb{A} \multimap \mathbb{A}$ is nondegenerate if and only if

ND1 any commutative square of solid arrows



can be filled in by dotted arrows to a commutative diagram as shown, and

ND2 any fork $b \xrightarrow{m} a \xrightleftharpoons[f']{f} c$ can be extended to a diagram



in which the triangle commutes and the right-hand column is a fork.

3 Coalgebras

We still need to prove that for any self-similarity system $(\mathbb{A}, M)$, the endofunctor $M \otimes -$ of $[\mathbb{A}, \mathbf{Set}]$ restricts to an endofunctor of $[\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$, and similarly with **Top** in place of **Set**. The set-theoretic case is straightforward.

Proposition 3.1 *Let $\mathbb{A}$ be a small category and $M : \mathbb{A} \multimap \mathbb{A}$ a nondegenerate module. Then*

$$M \otimes - : [\mathbb{A}, \mathbf{Set}] \longrightarrow [\mathbb{A}, \mathbf{Set}]$$

preserves nondegeneracy.

Nondegeneracy of M is also a *necessary* condition for $M \otimes -$ to preserve non-degeneracy: consider representables.

Proof Let $X : \mathbb{A} \longrightarrow \mathbf{Set}$ be nondegenerate. Then for any finite connected limit $\int_i Y_i$ in $[\mathbb{A}^{\text{op}}, \mathbf{Set}]$,

$$\int_i (Y_i \otimes M \otimes X) \cong \left(\int_i (Y_i \otimes M) \right) \otimes X \cong \left(\int_i Y_i \right) \otimes M \otimes X,$$

the first isomorphism by nondegeneracy of X and the second by nondegeneracy of M . So $M \otimes X$ is nondegenerate. $\square$

The topological case requires some preparatory lemmas. The first concerns $\mathbf{Set}$ -valued functors and follows immediately from Lemma 2.4.

Lemma 3.2 (Equality in $M \otimes X$) *Let $\mathbb{A}$ be a small category, let $M : \mathbb{A} \dashrightarrow \mathbb{A}$, and let $X \in [\mathbb{A}, \mathbf{Set}]_{\text{nondegen}}$. Take module elements*

$$\begin{array}{ccc} b & & b' \\ & \searrow m & \swarrow m' \\ & a & \end{array}$$

and $x \in Xb$, $x' \in Xb'$. Then $m \otimes x = m' \otimes x' \in (M \otimes X)a$ if and only if there exist a commutative square

$$\begin{array}{ccc} & c & \\ f \swarrow & & \searrow f' \\ b & & b' \\ m \searrow & & \swarrow m' \\ & a & \end{array}$$

and an element $z \in Xc$ such that $fz = x$ and $f'z = x'$. $\square$

Lemma 3.3 (Closed quotient map) *Let $\mathbb{A}$ be a small category, $X : \mathbb{A} \longrightarrow \mathbf{Top}$ a nondegenerate functor, and $Y : \mathbb{A}^{\text{op}} \longrightarrow \mathbf{Set}$ a functor whose category of elements is finite. Then the quotient map*

$$q : \sum_a Y a \times X a \longrightarrow \int^a Y a \times X a = Y \otimes X$$

is closed.

Proof A subset of $Y \otimes X$ is closed just when its inverse image under q is closed, so we must show that if V is a closed subset of $\sum Y a \times X a$ then its saturation $[V] = q^{-1}qV$ is also closed. Given $a \in \mathbb{A}$ and $y \in Y a$, write $V_{a,y}$ for the intersection of V with the (a,y) -summand $X a$ of

$$\sum_{(a,y) \in \mathbb{E}(Y)} X a \cong \sum_{a \in \mathbb{A}} Y a \times X a.$$

Then $[V] = \bigcup_{(a,y) \in \mathbb{E}(Y)} [V_{a,y}]$, so by finiteness of $\mathbb{E}(Y)$ it suffices to show that each $[V_{a,y}]$ is closed.

Fix $(a, y) \in \mathbb{E}(Y)$. By definition,

$$[V_{a,y}] = \{(a', y', x') \in \sum_{a' \in \mathbb{A}} Y a' \times X a' \mid y' \otimes x' = y \otimes x \text{ for some } x \in V_{a,y}\}.$$

So by nondegeneracy of X and Lemma 3.2, $(a', y', x') \in [V_{a,y}]$ if and only if:

there exist a span

$$\begin{array}{ccc} & b & \\ f \swarrow & & \searrow f' \\ a & & a' \end{array}$$

in $\mathbb{A}$ and $z \in Xb$ such that $fz \in V_{a,y}$, $f'z = x'$, and $yf = y'f'$,

or equivalently:

there exist a span

$$\begin{array}{ccc} & (b, w) & \\ f \swarrow & & \searrow f' \\ (a, y) & & (a', y') \end{array} \tag{13}$$

in $\mathbb{E}(Y)$ and $z \in Xb$ such that $fz \in V_{a,y}$ and $f'z = x'$.

So

$$[V_{a,y}] = \bigcup_{\text{spans (13)}} \{(a', y', x') \mid x' \in (Xf')(Xf)^{-1}V_{a,y}\}.$$

But each Xf is continuous and each Xf' closed, so each of the sets $\{\dots\}$ in this union is a closed subset of the (a', y') -summand Xa' . Moreover, finiteness of $\mathbb{E}(Y)$ guarantees that there are only finitely many spans of the form (13). So $[V_{a,y}]$ is a finite union of closed sets, hence closed. $\square$

Lemma 3.4 (Change of category) *Let $\mathcal{E}$ and $\mathcal{E}'$ be categories with finite colimits, $F : \mathcal{E} \longrightarrow \mathcal{E}'$ a functor preserving finite colimits, $\mathbb{A}$ a small category, and $M : \mathbb{A} \dashrightarrow \mathbb{A}$ a finite module. Then the square*

$$\begin{array}{ccc} [\mathbb{A}, \mathcal{E}] & \xrightarrow{M \otimes -} & [\mathbb{A}, \mathcal{E}] \\ F \circ - \downarrow & & \downarrow F \circ - \\ [\mathbb{A}, \mathcal{E}'] & \xrightarrow{M \otimes -} & [\mathbb{A}, \mathcal{E}'], \end{array}$$

commutes up to canonical isomorphism.

Proof Straightforward. $\square$

Proposition 3.5 *For all self-similarity systems $(\mathbb{A}, M)$, the functor*

$$M \otimes - : [\mathbb{A}, \mathbf{Top}] \longrightarrow [\mathbb{A}, \mathbf{Top}]$$

preserves nondegeneracy.

Proof Let $X \in [\mathbb{A}, \mathbf{Top}]_{\text{nondegen}}$. The functor $U : \mathbf{Top} \longrightarrow \mathbf{Set}$ preserves colimits (being left adjoint to the indiscrete space functor), so $M \otimes (U \circ X) \cong U \circ (M \otimes X)$ by Lemma 3.4. But $M \otimes (U \circ X)$ is nondegenerate by Proposition 3.1, so $U \circ (M \otimes X)$ is nondegenerate.

Now let $a \xrightarrow{f} a'$ be a map in $\mathbb{A}$, and consider the commutative square

$$\begin{array}{ccc} \sum_b M(b, a) \times Xb & \xrightarrow{\sum f_* \times 1} & \sum_b M(b, a') \times Xb \\ q_a \downarrow & & \downarrow q_{a'} \\ (M \otimes X)a & \xrightarrow{(M \otimes X)f} & (M \otimes X)a'. \end{array}$$

The map $\sum f_* \times 1$ is closed because each set $M(b, a)$ is finite. The map $q_{a'}$ is closed by Lemma 3.3 and finiteness of M . So $((M \otimes X)f) \circ q_a$ is closed; but q_a is a continuous surjection, so $(M \otimes X)f$ is closed. $\square$

Any M -coalgebra (X, ξ) in $\mathbf{Top}$ has an underlying M -coalgebra in $\mathbf{Set}$, since $U \circ X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate and there is a natural transformation

$$U\xi : U \circ X \longrightarrow U \circ (M \otimes X) \cong M \otimes (U \circ X).$$

This defines a functor

$$U_* : \mathbf{Coalg}(M, \mathbf{Top}) \longrightarrow \mathbf{Coalg}(M, \mathbf{Set})$$

where the domain and codomain are the categories of M -coalgebras in $\mathbf{Top}$ and $\mathbf{Set}$ respectively.

Proposition 3.6 (Top vs Set) *Let $(\mathbb{A}, M)$ be a self-similarity system. The forgetful functor*

$$U_* : \mathbf{Coalg}(M, \mathbf{Top}) \longrightarrow \mathbf{Coalg}(M, \mathbf{Set})$$

has a left adjoint, and if (I, ι) is a universal solution in $\mathbf{Top}$ then $U_(I, \iota)$ is a universal solution in $\mathbf{Set}$.*

Conversely, we will see that any universal solution in $\mathbf{Set}$ can be equipped with a topology that makes it the universal solution in $\mathbf{Top}$.

Proof Let D be the left adjoint to $U : \mathbf{Top} \longrightarrow \mathbf{Set}$, assigning to each set the corresponding discrete space. Then D preserves colimits, so commutes with $M \otimes -$ (Lemma 3.4). Moreover, if $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate then so is $D \circ X : \mathbb{A} \longrightarrow \mathbf{Top}$. Hence D induces a functor

$$D_* : \mathbf{Coalg}(M, \mathbf{Set}) \longrightarrow \mathbf{Coalg}(M, \mathbf{Top}).$$

For purely formal reasons, the adjunction $D \dashv U$ induces an adjunction $D_* \dashv U_*$. The statement on universal solutions follows from the fact that right adjoints preserve terminal objects. $\square$

4 The universal solution

In this section I construct the universal solution in **Set** and in **Top** of any given self-similarity system, assuming that the system satisfies a certain solvability condition **S**. In the Appendix I show that this sufficient condition **S** is also necessary. The construction therefore gives the universal solution whenever one exists.

Condition **S** on a self-similarity system $(\mathbb{A}, M)$ is:

S1 given any commutative diagram

$$\begin{array}{ccccccc}
 & & m_3 & & m_2 & & m_1 \\
 \cdots & \multimap & a_2 & \multimap & a_1 & \multimap & a_0 \\
 & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 \\
 \cdots & \multimap & b_2 & \multimap & b_1 & \multimap & b_0 \\
 & & \uparrow f'_2 & & \uparrow f'_1 & & \uparrow f'_0 \\
 \cdots & \multimap & a'_2 & \multimap & a'_1 & \multimap & a'_0, \\
 & & m'_3 & & m'_2 & & m'_1
 \end{array}$$

there exists a commutative square

$$\begin{array}{ccc}
 & a_0 & \\
 \nearrow & & \searrow f_0 \\
 \cdot & & b_0 \\
 \searrow & & \nearrow f'_0 \\
 & a'_0 &
 \end{array}$$

in $\mathbb{A}$, and

S2 given any serially commutative diagram

$$\begin{array}{ccccccc}
 & & m_3 & & m_2 & & m_1 \\
 \cdots & \multimap & a_2 & \multimap & a_1 & \multimap & a_0 \\
 & & \downarrow f_2 & \downarrow f'_2 & \downarrow f_1 & \downarrow f'_1 & \downarrow f_0 & \downarrow f'_0 \\
 \cdots & \multimap & b_2 & \multimap & b_1 & \multimap & b_0, \\
 & & p_3 & & p_2 & & p_1
 \end{array}$$

there exists a fork $\cdot \longrightarrow a_0 \begin{smallmatrix} \xrightarrow{f_0} \\ \xrightarrow{f'_0} \end{smallmatrix} b_0$ in $\mathbb{A}$.

Example 4.1 For any small category $\mathbb{A}$ there is a module $M : \mathbb{A} \multimap \mathbb{A}$ defined by $M(b, a) = \mathbb{A}(b, a)$, and $(\mathbb{A}, M)$ is a self-similarity system as long as $\sum_b \mathbb{A}(b, a)$ is finite for each $a \in \mathbb{A}$. Condition **S** says that $\mathbb{A}$ is componentwise cofiltered; so, for instance, the self-similarity system obtained by taking $\mathbb{A} = (0 \rightrightarrows 1)$ has no universal solution. If $\mathbb{A}$ is componentwise cofiltered then the universal solution is the functor $\mathbb{A} \longrightarrow \mathbf{Top}$ constant at the one-point space, with its unique coalgebra structure.

Here is the construction of the universal solution. The proofs that it works are in §5 (for **Set**) and §6 (for **Top**).

Let $(\mathbb{A}, M)$ be a self-similarity system. For each $a \in \mathbb{A}$, there is a category $\mathcal{J}a$ in which an object is an infinite sequence

$$\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0 = a \quad (14)$$

and a map $(a_\bullet, m_\bullet) \longrightarrow (a'_\bullet, m'_\bullet)$ is a commutative diagram

$$\begin{array}{ccccccc} \cdots & \xrightarrow{m_3} & a_2 & \xrightarrow{m_2} & a_1 & \xrightarrow{m_1} & a_0 = a \\ & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 = 1_a \\ \cdots & \xrightarrow{m'_3} & a'_2 & \xrightarrow{m'_2} & a'_1 & \xrightarrow{m'_1} & a'_0 = a. \end{array} \quad (15)$$

Moreover, each map $f : a \longrightarrow a'$ in $\mathbb{A}$ induces a functor $\mathcal{J}f : \mathcal{J}a \longrightarrow \mathcal{J}a'$, sending an object (14) of $\mathcal{J}a$ to the object

$$\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1 \xrightarrow{f m_1} a'$$

of $\mathcal{J}a'$. This defines a functor $\mathcal{J} : \mathbb{A} \longrightarrow \mathbf{Cat}$.

Write $\Pi_0 : \mathbf{Cat} \longrightarrow \mathbf{Set}$ for the functor sending a small category to its set of connected-components, and put $I = \Pi_0 \mathcal{J} : \mathbb{A} \longrightarrow \mathbf{Set}$; thus, Ia is a set of equivalence classes of diagrams (14). In §5 we will see that if $(\mathbb{A}, M)$ satisfies **S** then I is nondegenerate.

Warning 4.2 Ia is *not* the limit of finite approximations. Precisely, let $\mathcal{J}_n a$ be the category whose objects are diagrams of the form

$$a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0 = a \quad (16)$$

and whose arrows are commutative diagrams, and let $I_n a$ be the set of connected-components of $\mathcal{J}_n a$: then $\mathcal{J}a$ is the limit of the $(\mathcal{J}_n a)$ s, but Ia is not in general the limit of the $(I_n a)$ s. See the last paragraph of this section for an example.

There is an M -coalgebra structure ι on I defined by

$$\iota_a [\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] = (a_1 \xrightarrow{m_1} a) \otimes [\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1]$$

where $[]$ denotes connected-component. To see that this is a valid definition, note that given a map (15) in $\mathcal{J}a$, there is an equality $m_1 = m'_1 f_1$ and a map

$$\begin{array}{ccc} \cdots & \xrightarrow{m_3} & a_2 \xrightarrow{f_1 m_2} a'_1 \\ & & \downarrow f_2 \quad \downarrow 1 \\ \cdots & \xrightarrow{m'_3} & a'_2 \xrightarrow{m'_2} a'_1 \end{array}$$

in $\mathcal{I}a'_1$, so

$$\begin{aligned} m_1 \otimes [\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1] &= m'_1 \otimes [\cdots \xrightarrow{m_3} a_2 \xrightarrow{f_1 m_2} a'_1] \\ &= m'_1 \otimes [\cdots \xrightarrow{m'_3} a'_2 \xrightarrow{m'_2} a'_1]. \end{aligned}$$

Naturality of ι is easily checked. So if **S** holds then (I, ι) is an M -coalgebra, and it is in fact the terminal one, that is, the universal solution in **Set** (Theorem 5.11).

Now we construct the topology. For each $a \in \mathbb{A}$, $n \in \mathbb{N}$, and diagram of the form (16), there is a subset of Ia consisting of all those t such that

$$t = [\cdots \xrightarrow{m_{n+2}} a_{n+1} \xrightarrow{m_{n+1}} a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0 = a]$$

for some $m_{n+1}, a_{n+1}, m_{n+2}, \dots$. Generate a topology on Ia by taking each such subset to be closed. It is not obvious that the maps $If : Ia \longrightarrow Ia'$ are continuous or closed, or that the maps $\iota_a : Ia \longrightarrow (M \otimes I)a$ are continuous. Nevertheless, if **S** holds then they are, and so (I, ι) is an M -coalgebra in **Top**. Theorem 6.16 says that it is in fact the universal solution in **Top**.

(It is not hard to see that each of these basic closed subsets must be closed in *any* topology on (I, ι) [Lei2, 1.1]. So this is the coarsest possible topology.)

Let us see how all of this works in the case of the Freyd self-similarity system $(\mathbb{A}, M)$.

First, condition **S** holds. For **S1**, the only diagram

$$\begin{array}{ccc} a_0 & & \\ & \searrow f_0 & \\ & & b_0 \\ & \nearrow f'_0 & \\ a'_0 & & \end{array}$$

in $\mathbb{A}$ that cannot be completed to a commutative square is (up to symmetry) that in which $f_0 = \sigma$ and $f'_0 = \tau$, and then there is no infinite commutative diagram as in **S1**: indeed, there is not even a commutative diagram of the form

$$\begin{array}{ccc} \cdot & \xrightarrow{\quad} & 0 \\ \downarrow & & \downarrow \sigma \\ \cdot & \xrightarrow{\quad} & 1 \\ \uparrow & & \uparrow \tau \\ \cdot & \xrightarrow{\quad} & 0. \end{array}$$

Similarly, for **S2**, the only parallel pair of arrows in $\mathbb{A}$ that cannot be completed

to a fork is $0 \xrightarrow[\tau]{\sigma} 1$, and there is no serially commutative diagram of the form

$$\begin{array}{ccc} \cdot & \xrightarrow{\quad} & 0 \\ \downarrow & & \downarrow \sigma \\ \cdot & \xrightarrow{\quad} & 1. \end{array}$$

The universal solution (I, ι) has $I1 = [0, 1]$, so according to the construction, an element of $[0, 1]$ is an equivalence class of diagrams

$$\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1 \xrightarrow{m_1} 1.$$

If each a_n is 1 then each m_n is either $[0, \frac{1}{2}]$ or $[\frac{1}{2}, 1]$ and the diagram is just a binary expansion: for instance,

$$\cdots \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[0, \frac{1}{2}]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[0, \frac{1}{2}]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1$$

corresponds to $0.10101 \dots$, representing $\frac{2}{3} \in [0, 1]$. Otherwise, the diagram is of the form

$$\cdots \xrightarrow{\text{id}} 0 \xrightarrow{\text{id}} 0 \xrightarrow{m_{n+1}} 1 \xrightarrow{m_n} \cdots \xrightarrow{m_1} 1$$

where $m_1, \dots, m_n \in \{[0, \frac{1}{2}], [\frac{1}{2}, 1]\}$ and $m_{n+1} \in \{0, \frac{1}{2}, 1\}$: for instance,

$$\cdots \xrightarrow{\text{id}} 0 \xrightarrow{\text{id}} 0 \xrightarrow{\frac{1}{2}} 1 \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[0, \frac{1}{2}]} 1.$$

To see which element t of $[0, 1]$ this represents, reason as follows: the $[0, \frac{1}{2}]$ says that $t \in [0, \frac{1}{2}]$; the right-hand copy of $[\frac{1}{2}, 1]$ says that t is in the upper half of $[0, \frac{1}{2}]$, that is, in $[\frac{1}{4}, \frac{1}{2}]$; the left-hand copy of $[\frac{1}{2}, 1]$ says that t is in the upper half of $[\frac{1}{4}, \frac{1}{2}]$, that is, in $[\frac{3}{8}, \frac{1}{2}]$; then the $\frac{1}{2}$ says that t is the midpoint of $[\frac{3}{8}, \frac{1}{2}]$, that is, $t = \frac{7}{16}$.

An element of $[0, 1]$ has at most two binary expansions, but may have infinitely many representations in $\mathbb{J}1$. For instance, the representations of $\frac{1}{2}$ are

$$\cdots \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[0, \frac{1}{2}]} 1, \quad (17)$$

$$\cdots \xrightarrow{[0, \frac{1}{2}]} 1 \xrightarrow{[0, \frac{1}{2}]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1, \quad (18)$$

$$\cdots \xrightarrow{\text{id}} 0 \xrightarrow{\text{id}} 0 \xrightarrow{\frac{1}{2}} 1, \quad (19)$$

and for any $n \in \mathbb{N}$,

$$\cdots \xrightarrow{\text{id}} 0 \xrightarrow{1} 1 \xrightarrow{[\frac{1}{2}, 1]} \cdots \xrightarrow{[\frac{1}{2}, 1]} 1 \xrightarrow{[0, \frac{1}{2}]} 1, \quad (20)$$

$$\cdots \xrightarrow{\text{id}} 0 \xrightarrow{0} 1 \xrightarrow{[0, \frac{1}{2}]} \cdots \xrightarrow{[0, \frac{1}{2}]} 1 \xrightarrow{[\frac{1}{2}, 1]} 1 \quad (21)$$

with n copies of $[\frac{1}{2}, 1]$ and $[0, \frac{1}{2}]$ respectively.

The construction says that two objects of $\mathcal{I}1$ represent the same element of $[0, 1]$ if and only if they are in the same connected-component. So, for instance, each of (17)–(21) should be in the same component; the connected diagram

$$\begin{array}{ccccccc}
 & \xrightarrow{[\frac{1}{2}, 1]} & 1 & \xrightarrow{[\frac{1}{2}, 1]} & 1 & \xrightarrow{[0, \frac{1}{2}]} & 1 \\
 & \uparrow \text{id} & \uparrow \tau & \uparrow \text{id} & \uparrow \tau & \uparrow \frac{1}{2} & \uparrow 1 \\
 \cdots & \xrightarrow{\quad} & 0 & \xrightarrow{\quad} & 0 & \xrightarrow{\quad} & 1 \\
 & \downarrow \sigma & \downarrow \sigma & \downarrow \sigma & \downarrow \sigma & \downarrow 1 & \downarrow 1 \\
 & \xrightarrow{[0, \frac{1}{2}]} & 1 & \xrightarrow{[0, \frac{1}{2}]} & 1 & \xrightarrow{[\frac{1}{2}, 1]} & 1 \\
 & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow
 \end{array}$$

shows that (17)–(19) are, and the others are left to the reader. Observe in general that any two objects of $\mathcal{I}a$ of the form

$$\cdots \xrightarrow{m'_{n+2}} a'_{n+1} \xrightarrow{m'_{n+1}} a'_n \xrightarrow{m_n f} a_{n-1} \xrightarrow{m_{n-1}} \cdots \xrightarrow{m_1} a_0 = a, \quad (22)$$

$$\cdots \xrightarrow{m'_{n+2}} a'_{n+1} \xrightarrow{f m'_{n+1}} a_n \xrightarrow{m_n} a_{n-1} \xrightarrow{m_{n-1}} \cdots \xrightarrow{m_1} a_0 = a \quad (23)$$

(where $f : a'_n \longrightarrow a_n$) are in the same connected-component, because there is a map

$$(\dots, 1_{a'_{n+1}}, f, 1_{a_{n-1}}, \dots, 1_{a_0})$$

from the top row to the bottom.

The constructed topology on $[0, 1]$ is generated by taking as closed all subsets of the form $[k/2^n, l/2^n]$ where $k, l, n \in \mathbb{N}$ and $l \in \{k, k+1\}$. This is exactly the Euclidean topology.

Finally, this example shows that Ia need not be the limit of finite approximations (Warning 4.2). It is not hard to show that for each n , the category $\mathcal{I}_n 1$ is connected: so each $I_n 1$ is a one-element set, and $I1 = [0, 1]$ is plainly not the limit of the $(I_n 1)$ s.

5 Set-theoretic proofs

Fix a self-similarity system $(\mathbb{A}, M)$ satisfying **S**. In this section I prove that the functor $I : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate and that (I, ι) is the universal solution.

Nondegeneracy of I will follow from a kind of nondegeneracy property of $\mathcal{I} : \mathbb{A} \longrightarrow \mathbf{Cat}$. Any **Cat**-valued functor $\mathcal{X} : \mathbb{B} \longrightarrow \mathbf{Cat}$ has a **category of elements** $\mathbb{E}(\mathcal{X})$, in which an object is a pair (b, x) with $b \in \mathbb{B}$ and $x \in \mathcal{X}b$ and an arrow $(b, x) \longrightarrow (b', x')$ is a pair (g, ξ) with $g : b \longrightarrow b'$ in $\mathbb{B}$ and $\xi : (\mathcal{X}g)x \longrightarrow x'$ in $\mathcal{X}b'$. This is related to the notion of the category of elements of a **Set**-valued functor $X : \mathbb{B} \longrightarrow \mathbf{Set}$ (page 12) by the isomorphism $\mathbb{E}(X) \cong \mathbb{E}(D \circ X)$, where $D : \mathbf{Set} \longrightarrow \mathbf{Cat}$ is the functor assigning to each set

the corresponding discrete category. The nondegeneracy property mentioned is that $\mathbb{E}(\mathcal{J})$ is componentwise cofiltered (that is, $\mathbb{E}(\mathcal{J})^{\text{op}}$ is componentwise filtered, in the sense defined after Corollary 2.7).

This is proved by a finiteness argument. Notation: if $\mathcal{L}$ is the limit of a diagram

$$\cdots \longrightarrow \mathcal{L}_3 \longrightarrow \mathcal{L}_2 \longrightarrow \mathcal{L}_1$$

in some category, I write pr_n for both the projection $\mathcal{L} \longrightarrow \mathcal{L}_n$ and the given map $\mathcal{L}_m \longrightarrow \mathcal{L}_n$ for any $m \geq n$.

Lemma 5.1 (König [Kön]) *The limit in **Set** of a diagram*

$$\cdots \longrightarrow \mathcal{F}_3 \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F}_1$$

of finite nonempty sets is nonempty. More precisely, for any sequence $(F_n)_{n \geq 1}$ with $F_n \in \mathcal{F}_n$ there exists an element G of the limit such that

$$\forall r \geq 1, \exists n \geq r : \text{pr}_r(F_n) = \text{pr}_r(G).$$

Remark The first sentence is a special case of the fact that a componentwise cofiltered limit of nonempty compact Hausdorff spaces is nonempty (compare [Bou, I.9.6]).

Proof Take a sequence $(F_n)_{n \geq 1}$ with $F_n \in \mathcal{F}_n$. We define, for each $r \geq 1$, an infinite subset $\mathbb{N}_r$ of $\mathbb{N}$ and an element $G_r \in \mathcal{F}_r$ such that

- for all $r \geq 1$, $\mathbb{N}_r \subseteq \mathbb{N}_{r-1} \cap \{r, r+1, \dots\}$ (writing $\mathbb{N}_0 = \mathbb{N}$)
- for all $r \geq 1$ and $n \in \mathbb{N}_r$, $\text{pr}_r(F_n) = G_r$.

Suppose inductively that $r \geq 1$ and $\mathbb{N}_{r-1}$ is defined. As n runs through the infinite set $\mathbb{N}_{r-1} \cap \{r, r+1, \dots\}$, $\text{pr}_r(F_n)$ takes values in the finite set $\mathcal{F}_r$, so takes some value $G_r \in \mathcal{F}_r$ infinitely often. Putting

$$\mathbb{N}_r = \{n \in \mathbb{N}_{r-1} \cap \{r, r+1, \dots\} \mid \text{pr}_r(F_n) = G_r\}$$

completes the induction.

For each $r \geq 1$ we have $G_r = \text{pr}_r(G_{r+1})$, since we may choose $n \in \mathbb{N}_{r+1}$ and then

$$\text{pr}_r(G_{r+1}) = \text{pr}_r(\text{pr}_{r+1}(F_n)) = \text{pr}_r(F_n) = G_r.$$

So there is a unique element G of the limit such that $\text{pr}_r(G) = G_r$ for all $r \geq 1$. Given $r \geq 1$, we may choose $n \in \mathbb{N}_r$, and then $n \geq r$ and $\text{pr}_r(F_n) = G_r = \text{pr}_r(G)$ as required. $\square$

This will be applied as follows. Suppose we have a limit $\mathcal{L}$ of categories

$$\cdots \longrightarrow \mathcal{L}_3 \longrightarrow \mathcal{L}_2 \longrightarrow \mathcal{L}_1,$$

categories $\mathbb{J} \subseteq \mathbb{K}$, and a diagram D of shape $\mathbb{J}$ in $\mathcal{L}$, and suppose we are interested in extending D to a diagram of shape $\mathbb{K}$. If we can do so then we can certainly

extend $\text{pr}_n \circ D : \mathbb{J} \longrightarrow \mathcal{L}_n$ to $\mathbb{K}$ for all n ; the converse does not hold, because we cannot necessarily choose the extensions $\mathbb{K} \longrightarrow \mathcal{L}_n$ in a coherent way. But the following lemma says that we can do it if for each n there are only finitely many choices of extension.

Lemma 5.2 (Factorization) *Let $\mathcal{L}$ be the limit of a diagram*

$$\cdots \longrightarrow \mathcal{L}_3 \longrightarrow \mathcal{L}_2 \longrightarrow \mathcal{L}_1$$

in some category $\mathcal{E}$. Let $P : \mathbb{J} \longrightarrow \mathbb{K}$ and $D : \mathbb{J} \longrightarrow \mathcal{L}$ be maps in $\mathcal{E}$ such that for each $n \geq 1$, the set of factorizations of $\text{pr}_n \circ D$ through P is finite and nonempty. Then D factors through P .

Proof Apply Lemma 5.1 with $\mathcal{F}_n = \{F \in \mathcal{E}(\mathbb{K}, \mathcal{L}_n) \mid F \circ P = \text{pr}_n \circ D\}$. $\square$

For each $n \in \mathbb{N}$ we have a functor $\mathcal{I}_n : \mathbb{A} \longrightarrow \mathbf{Cat}$ (see 4.2). The evident projections make $\mathbb{E}(\mathcal{I})$ the limit in $\mathbf{Cat}$ of

$$\cdots \longrightarrow \mathbb{E}(\mathcal{I}_2) \longrightarrow \mathbb{E}(\mathcal{I}_1).$$

Proposition 5.3 $\mathbb{E}(\mathcal{I})$ is componentwise cofiltered.

Proof We have to prove that every diagram $\cdot \longrightarrow \cdot \longleftarrow \cdot$ in $\mathbb{E}(\mathcal{I})$ can be completed to a commutative square and that every parallel pair $\cdot \rightrightarrows \cdot$ can be completed to a fork. The two cases are very similar, so I just do the first.

Take a diagram

$$\begin{array}{ccccccc} & & m_3 & & m_2 & & m_1 \\ & & \dashrightarrow & & \dashrightarrow & & \dashrightarrow \\ \cdots & & a_2 & & a_1 & & a_0 \\ & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 \\ \cdots & & b_2 & & b_1 & & b_0 \\ & & \uparrow f'_2 & & \uparrow f'_1 & & \uparrow f'_0 \\ \cdots & & a'_2 & & a'_1 & & a'_0 \\ & & \dashleftarrow m'_3 & & \dashleftarrow m'_2 & & \dashleftarrow m'_1 \end{array} \quad (24)$$

of shape $(\cdot \longrightarrow \cdot \longleftarrow \cdot)$ in $\mathbb{E}(\mathcal{I})$. We apply Lemma 5.2 where P is the inclusion

$$\mathbb{J} = \left(\begin{array}{ccc} & & \\ \cdot & & \cdot \\ & \searrow & \swarrow \\ & \cdot & \end{array} \right) \hookrightarrow \left(\begin{array}{ccc} & \cdot & \\ \swarrow & & \searrow \\ \cdot & & \cdot \\ \searrow & & \swarrow \\ & \cdot & \end{array} \right) = \mathbb{K}$$

(the square in $\mathbb{K}$ being commutative), $\mathcal{L}_n = \mathbb{E}(\mathcal{I}_n)$, $\mathcal{L} = \mathbb{E}(\mathcal{I})$, and D is the diagram (24) with its rightmost block removed. The hypothesis of Lemma 5.2

is that for each $n \geq 1$, the set of diagrams of the form

$$\begin{array}{ccccccc}
 a_n & \xrightarrow{m_n} & a_{n-1} & \xrightarrow{m_{n-1}} & \cdots & \xrightarrow{m_2} & a_1 \\
 g_n \uparrow & & g_{n-1} \uparrow & & & & g_1 \uparrow \\
 c_n & \xrightarrow{q_n} & c_{n-1} & \xrightarrow{q_{n-1}} & \cdots & \xrightarrow{q_2} & c_1 \\
 g'_n \downarrow & & g'_{n-1} \downarrow & & & & g'_1 \downarrow \\
 a'_n & \xrightarrow{m'_n} & a'_{n-1} & \xrightarrow{m'_{n-1}} & \cdots & \xrightarrow{m'_2} & a'_1
 \end{array} \tag{25}$$

satisfying $f_1 g_1 = f'_1 g'_1, \dots, f_n g_n = f'_n g'_n$ is nonempty and finite.

Finiteness follows from finiteness of M (and the absence of the rightmost block). For nonemptiness, let $n \geq 1$. Then **S1** implies that there exist c_n , g_n , and g'_n making

$$\begin{array}{ccccc}
 & a_n & \xrightarrow{m_n} & a_{n-1} & \\
 g_n \nearrow & & & & \searrow f_{n-1} \\
 c_n & & & b_n & \xrightarrow{p_n} b_{n-1} \\
 g'_n \searrow & & f'_n \nearrow & & \nearrow f'_{n-1} \\
 & a'_n & \xrightarrow{m'_n} & a'_{n-1} &
 \end{array}$$

commute, and then nondegeneracy of M (condition **ND1** at the end of §2) implies that the outside of this diagram can also be filled in as

$$\begin{array}{ccccc}
 & a_n & \xrightarrow{m_n} & a_{n-1} & \\
 g_n \nearrow & & & g_{n-1} \nearrow & \searrow f_{n-1} \\
 c_n & \xrightarrow{q_n} & c_{n-1} & & b_{n-1} \\
 g'_n \searrow & & g'_{n-1} \searrow & & \nearrow f'_{n-1} \\
 & a'_n & \xrightarrow{m'_n} & a'_{n-1} &
 \end{array}$$

Repeating this argument $(n-2)$ times gives a diagram (25), as required.

So the hypothesis of Lemma 5.2 holds, and D can be completed to a commutative square in $\mathbb{E}(\mathcal{J})$. Using the diagram-filling argument one more time shows that (24) can be, too. $\square$

The next few results show that for general reasons, $\mathbb{E}(\mathcal{J})$ being component-wise cofiltered implies that each $\mathcal{J}a$ is too and that $I : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate.

Lemma 5.4 *Let $\mathcal{J} : \mathbb{B} \longrightarrow \mathbf{Cat}$ be a functor on a small category $\mathbb{B}$. If $\mathbb{E}(\mathcal{J})$ is componentwise cofiltered then $\mathcal{J}a$ is componentwise cofiltered for each $a \in \mathbb{B}$.*

Proof We have to prove that every diagram $\cdot \longrightarrow \cdot \longleftarrow \cdot$ in $\mathcal{J}a$ can be completed to a commutative square and that every parallel pair $\cdot \rightrightarrows \cdot$ can be completed to a fork. Again I just do the first case; the second is similar.

Take a diagram

$$\begin{array}{ccc} \omega & & \omega' \\ & \searrow \phi & \swarrow \phi' \\ & \chi & \end{array}$$

in $\mathcal{J}a$. Then there is a commutative square

$$\begin{array}{ccc} & (b, \zeta) & \\ (g, \gamma) \swarrow & & \searrow (g', \gamma') \\ (a, \omega) & & (a, \omega') \\ (1, \phi) \searrow & & \swarrow (1, \phi') \\ & (a, \chi) & \end{array}$$

in $\mathbb{E}(\mathcal{J})$. Commutativity says that $g = g'$ and that the square

$$\begin{array}{ccc} & (\mathcal{J}g)\zeta & \\ \gamma \swarrow & & \searrow \gamma' \\ \omega & & \omega' \\ \phi \searrow & & \swarrow \phi' \\ & \chi & \end{array}$$

in $\mathcal{J}a$ commutes, as required. $\square$

Proposition 5.5 *$\mathcal{J}a$ is componentwise cofiltered for each $a \in \mathbb{A}$.* $\square$

Lemma 5.6 *Let $\mathcal{J} : \mathbb{B} \longrightarrow \mathbf{Cat}$ be a functor on a small category $\mathbb{B}$. If $\mathbb{E}(\mathcal{J})$ is componentwise cofiltered then so is $\mathbb{E}(\Pi_0 \mathcal{J})$.*

Proof Once again the proof splits into two similar cases. For variety, I do the second: that every diagram $(a, [\omega]) \xrightleftharpoons[f']{f} (b, [\chi])$ in $\mathbb{E}(\Pi_0 \mathcal{J})$ extends to a fork.

Since $[(\mathcal{J}f)\omega] = [\chi] = [(\mathcal{J}f')\omega]$, Lemmas 2.3 and 5.4 imply that there exists a span

$$\begin{array}{ccc} & \xi & \\ \delta \swarrow & & \searrow \delta' \\ (\mathcal{J}f)\omega & & (\mathcal{J}f')\omega \end{array}$$

in $\mathcal{J}b$. We therefore have a finite connected diagram (solid arrows)

$$\begin{array}{ccc}
 & (c, \zeta) & \\
 (g, \gamma) \swarrow \cdots & & \searrow \cdots (fg=f'g, \theta) \\
 (a, \omega) & & (b, \xi) \\
 \downarrow (f, 1) & \swarrow (f', 1) \quad \searrow (1, \delta) & \downarrow (1, \delta') \\
 (b, (\mathcal{J}f)\omega) & & (b, (\mathcal{J}f')\omega)
 \end{array}$$

in $\mathbb{E}(\mathcal{J})$, so by hypothesis there exists a dotted commutative diagram, giving a fork

$$(c, [\zeta]) \xrightarrow{g} (a, [\omega]) \xrightleftharpoons[f']{f} (b, [\chi])$$

in $\mathbb{E}(\Pi_0\mathcal{J})$. □

Proposition 5.7 (*I nondegenerate*) $I : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate. □

Hence (I, ι) is an M -coalgebra. By Lambek's Lemma, a necessary condition for it to be the universal solution is that ι is an isomorphism, and we can prove this immediately.

Proposition 5.8 (*I is a fixed point*) $\iota : I \longrightarrow M \otimes I$ is an isomorphism.

Proof It is enough to show that $\iota_a : Ia \longrightarrow (M \otimes I)a$ is bijective for each $a \in \mathbb{A}$. Certainly ι_a is surjective. For injectivity, suppose that

$$\iota_a[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] = \iota_a[\cdots \xrightarrow{m'_2} a'_1 \xrightarrow{m'_1} a],$$

that is,

$$m_1 \otimes [\cdots \xrightarrow{m_2} a_1] = m'_1 \otimes [\cdots \xrightarrow{m'_2} a'_1].$$

By Lemma 3.2 and nondegeneracy of I , there exist a commutative square

$$\begin{array}{ccc}
 & b & \\
 f \swarrow & & \searrow f' \\
 a_1 & & a'_1 \\
 m_1 \searrow & & \swarrow m'_1 \\
 & a &
 \end{array}$$

and an element $[\cdots \xrightarrow{p_2} b_1 \xrightarrow{p_1} b] \in Ib$ such that

$$\begin{aligned}
 [\cdots \xrightarrow{p_2} b_1 \xrightarrow{fp_1} a_1] &= [\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1], \\
 [\cdots \xrightarrow{p_2} b_1 \xrightarrow{f'p_1} a'_1] &= [\cdots \xrightarrow{m'_3} a'_2 \xrightarrow{m'_2} a'_1].
 \end{aligned}$$

Then

$$\begin{aligned} [\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] &= [\cdots \xrightarrow{p_2} b_1 \xrightarrow{fp_1} a_1 \xrightarrow{m_1} a] \\ &= [\cdots \xrightarrow{p_2} b_1 \xrightarrow{p_1} b \xrightarrow{m_1 f} a], \end{aligned}$$

using the observation at (22) and (23) (page 28). But $m_1 f = m'_1 f'$, so by symmetry of argument,

$$[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] = [\cdots \xrightarrow{m'_2} a'_1 \xrightarrow{m'_1} a]$$

as required. $\square$

Here is the key concept for the rest of the proof. Let (X, ξ) be an M -coalgebra, $a \in \mathbb{A}$, and $x \in Xa$. A **resolution** of x is a diagram

$$\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0 = a \quad (26)$$

together with a sequence $(x_n)_{n \in \mathbb{N}}$ such that $x_n \in Xa_n$, $x_0 = x$, and

$$\xi(x_n) = m_{n+1} \otimes x_{n+1}$$

for all $n \in \mathbb{N}$. I will also call $(x_n)_{n \in \mathbb{N}}$ a **resolution of x along** the diagram (26). Clearly every element x of a coalgebra has at least one resolution.

Lemma 5.9 (Direction of resolution) *Let (X, ξ) be a nondegenerate M -coalgebra, $a \in \mathbb{A}$, and $x \in Xa$. If x has resolutions along both*

$$(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0 = a) \quad \text{and} \quad (\cdots \xrightarrow{m'_2} a'_1 \xrightarrow{m'_1} a'_0 = a)$$

then

$$[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0] = [\cdots \xrightarrow{m'_2} a'_1 \xrightarrow{m'_1} a'_0].$$

Proof Choose sequences $(x_n)_{n \in \mathbb{N}}$ and $(x'_n)_{n \in \mathbb{N}}$ resolving x along the two diagrams respectively. I construct by induction a commutative diagram

$$\begin{array}{ccccc} \cdots & \xrightarrow{m_3} & a_2 & \xrightarrow{m_2} & a_1 \xrightarrow{m_1} a_0 = a \\ & & \uparrow f_2 & & \uparrow f_1 \\ \cdots & \xrightarrow{p_3} & b_2 & \xrightarrow{p_2} & b_1 \xrightarrow{p_1} b_0 = a \\ & & \downarrow f'_2 & & \downarrow f'_1 \\ \cdots & \xrightarrow{m'_3} & a'_2 & \xrightarrow{m'_2} & a'_1 \xrightarrow{m'_1} a'_0 = a \\ & & & & \downarrow f'_0 = 1_a \end{array}$$

and a sequence $(y_n)_{n \in \mathbb{N}}$ such that $y_n \in Xb_n$, $f_n y_n = x_n$, and $f'_n y_n = x'_n$ for each $n \in \mathbb{N}$. For the base step, take $y_0 = x$. For the inductive step, let $n \in \mathbb{N}$ and suppose that b_n , f_n , f'_n , and y_n are constructed. We may write

$$\xi(y_n) = (c \xrightarrow{q} b_n) \otimes z$$

with $z \in Xc$. Then

$$m_{n+1} \otimes x_{n+1} = \xi(x_n) = \xi(f_n y_n) = f_n \xi(y_n) = (f_n q) \otimes z,$$

so by nondegeneracy of X and Lemma 3.2, there exist a commutative diagram as labelled (a) below and an element $w \in Xd$ such that $gw = z$ and $hw = x_{n+1}$:

$$\begin{array}{ccccc}
 & & a_{n+1} & \xrightarrow{m_{n+1}} & a_n \\
 & & \uparrow h & & \uparrow f_n \\
 & & d & \xrightarrow{g} & c \\
 b_{n+1} & \xrightarrow{k} & & & \\
 & & \downarrow k' & & \downarrow f'_n \\
 & & d' & \xrightarrow{g'} & c \\
 & & \downarrow h' & & \downarrow f'_n \\
 & & a'_{n+1} & \xrightarrow{m'_{n+1}} & a'_n
 \end{array}
 \quad \begin{array}{c} \text{(a)} \\ \text{(b)} \\ \text{(a')} \end{array}$$

Similarly, there exist a commutative diagram (a') and $w' \in Xd'$ such that $g'w' = z$ and $h'w' = x'_{n+1}$. So by nondegeneracy of X (condition **ND1**), there exist a commutative square (b) and $y_{n+1} \in Xb_{n+1}$ such that $ky_{n+1} = w$ and $k'y_{n+1} = w'$. Putting $p_{n+1} = qgk$, $f_{n+1} = hk$, and $f'_{n+1} = h'k'$ completes the induction. $\square$

Now consider resolutions in the coalgebra (I, ι) . Given any $a \in \mathbb{A}$ and

$$(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0) \in \mathcal{I}a, \quad (27)$$

there is a canonical resolution of $[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0] \in Ia$, consisting of (27) itself together with $[\cdots \xrightarrow{m_{n+2}} a_{n+1} \xrightarrow{m_{n+1}} a_n] \in Ia_n$ as ' x_n '.

Proposition 5.10 (Double complex) *Let*

$$\begin{array}{ccccccc}
 & & & & & & \vdots \\
 & & & & & & \downarrow m_3 \\
 & & & & & & \vdots \\
 & & & & & & \downarrow m_2 \\
 & & & & & & \vdots \\
 & & & & & & \downarrow m_1 \\
 & & & & & & \vdots \\
 \cdots & \xrightarrow{m_2^3} & a_2^2 & \xrightarrow{m_2^2} & a_2^1 & \xrightarrow{m_2^1} & a_2^0 \\
 & & & & & & \downarrow m_2 \\
 \cdots & \xrightarrow{m_1^3} & a_1^2 & \xrightarrow{m_1^2} & a_1^1 & \xrightarrow{m_1^1} & a_1^0 \\
 & & & & & & \downarrow m_1 \\
 \cdots & \xrightarrow{m_0^3} & a_0^2 & \xrightarrow{m_0^2} & a_0^1 & \xrightarrow{m_0^1} & a_0^0
 \end{array}$$

be a diagram satisfying

$$[\cdots \xrightarrow{m_n^3} a_n^2 \xrightarrow{m_n^2} a_n^1 \xrightarrow{m_n^1} a_n^0] = [\cdots \xrightarrow{m_{n+1}^2} a_{n+1}^1 \xrightarrow{m_{n+1}^1} a_{n+1}^0 \xrightarrow{m_{n+1}} a_n^0]$$

for all $n \in \mathbb{N}$. Then

$$[\cdots \xrightarrow{m_0^3} a_0^2 \xrightarrow{m_0^2} a_0^1 \xrightarrow{m_0^1} a_0^0] = [\cdots \xrightarrow{m_3} a_2^0 \xrightarrow{m_2} a_1^0 \xrightarrow{m_1} a_0^0]. \quad (28)$$

Proof The left-hand side of (28) can be resolved canonically along

$$\cdots \xrightarrow{m_0^2} a_0^1 \xrightarrow{m_0^1} a_0^0.$$

It also has a resolution $(x_n)_{n \in \mathbb{N}}$ along

$$\cdots \xrightarrow{m_2} a_1^0 \xrightarrow{m_1} a_0^0,$$

where

$$x_n = [\cdots \xrightarrow{m_n^2} a_n^1 \xrightarrow{m_n^1} a_n^0] \in Ia_n^0,$$

since by hypothesis

$$\begin{aligned}
 \iota x_n &= \iota [\cdots \xrightarrow{m_{n+1}^1} a_{n+1}^0 \xrightarrow{m_{n+1}} a_n^0] \\
 &= m_{n+1} \otimes x_{n+1}.
 \end{aligned}$$

The result follows from nondegeneracy of I and Lemma 5.9. $\square$

Theorem 5.11 (Universal solution in Set) (I, ι) is the universal solution of $(\mathbb{A}, M)$ in **Set**.

Proof Let (X, ξ) be a nondegenerate coalgebra. We have to show that there is a unique map $(X, \xi) \longrightarrow (I, \iota)$.

Existence Given any $a \in \mathbb{A}$ and $x \in Xa$, we may choose a resolution

$$((\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0), (x_0, x_1, x_2, \dots)) \quad (29)$$

of x and put

$$\bar{\xi}_a(x) = [\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] \in Ia.$$

This defines for each a a function $\bar{\xi}_a : Xa \longrightarrow Ia$, which by Lemma 5.9 is independent of choice of resolution. I claim that $\bar{\xi}$ is a map $(X, \xi) \longrightarrow (I, \iota)$ of coalgebras. First, it is a natural transformation, that is, if $a \xrightarrow{f} a'$ is a map in $\mathbb{A}$ and $x \in Xa$ then $\bar{\xi}_{a'}(fx) = f\bar{\xi}_a(x)$. For choose a resolution (29) of x : then

$$((\cdots \xrightarrow{m_2} a_1 \xrightarrow{fm_1} a'), (fx, x_1, x_2, \dots))$$

is a resolution of fx , so

$$\begin{aligned} \bar{\xi}_{a'}(fx) &= [\cdots \xrightarrow{m_2} a_1 \xrightarrow{fm_1} a'] \\ &= f[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] \\ &= f\bar{\xi}_a(x). \end{aligned}$$

Second, $\bar{\xi}$ is a map of coalgebras, that is, if $a \in \mathbb{A}$ and $x \in Xa$ then

$$(M \otimes \bar{\xi})_a \xi_a(x) = \iota_a \bar{\xi}_a(x).$$

For choose a resolution (29) of x : then

$$((\cdots \xrightarrow{m_3} a_2 \xrightarrow{m_2} a_1), (x_1, x_2, x_3, \dots))$$

is a resolution of x_1 , so

$$\begin{aligned} (M \otimes \bar{\xi})_a \xi_a(x) &= (M \otimes \bar{\xi})_a(m_1 \otimes x_1) \\ &= m_1 \otimes \bar{\xi}_{a_1}(x_1) \\ &= m_1 \otimes [\cdots \xrightarrow{m_2} a_1] \\ &= \iota_a[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a] \\ &= \iota_a \bar{\xi}_a(x). \end{aligned}$$

Uniqueness Let $\tilde{\xi} : (X, \xi) \longrightarrow (I, \iota)$ be a map of coalgebras, $a \in \mathbb{A}$, and $x \in Xa$. We show that $\tilde{\xi}_a(x) = \bar{\xi}_a(x)$.

Choose a resolution (29) of x , and for each $n \in \mathbb{N}$, write

$$\tilde{\xi}_{a_n}(x_n) = [\cdots \xrightarrow{m_n^2} a_n^1 \xrightarrow{m_n^1} a_n^0 = a_n].$$

For each $n \in \mathbb{N}$, we have

$$(M \otimes \tilde{\xi})_{a_n} \xi_{a_n}(x_n) = \iota_{a_n} \tilde{\xi}_{a_n}(x_n)$$

by definition of map of coalgebras. On the other hand,

$$\begin{aligned} (M \otimes \tilde{\xi})_{a_n} \xi_{a_n}(x_n) &= (M \otimes \tilde{\xi})_{a_n}(m_{n+1} \otimes x_{n+1}) \\ &= m_{n+1} \otimes \tilde{\xi}_{a_{n+1}}(x_{n+1}) \\ &= m_{n+1} \otimes [\cdots \xrightarrow{m_{n+1}^2} a_{n+1}^1 \xrightarrow{m_{n+1}^1} a_{n+1}^0 = a_{n+1}] \\ &= \iota_{a_n} [\cdots \xrightarrow{m_{n+1}^1} a_{n+1}^0 = a_{n+1} \xrightarrow{m_{n+1}} a_n]. \end{aligned}$$

Since ι_{a_n} is injective (Proposition 5.8),

$$\begin{aligned} [\cdots \xrightarrow{m_{n+1}^1} a_{n+1}^0 = a_{n+1} \xrightarrow{m_{n+1}} a_n] &= \tilde{\xi}_{a_n}(x_n) \\ &= [\cdots \xrightarrow{m_n^2} a_n^1 \xrightarrow{m_n^1} a_n^0 = a_n] \end{aligned}$$

for each $n \in \mathbb{N}$. So Proposition 5.10 applies, and

$$[\cdots \xrightarrow{m_0^2} a_0^1 \xrightarrow{m_0^1} a_0^0] = [\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0] \in Ia,$$

that is, $\tilde{\xi}_a(x) = \bar{\xi}_a(x)$, as required. $\square$

6 Topological proofs

Fix a self-similarity system $(\mathbb{A}, M)$. In this section I show that if $(\mathbb{A}, M)$ satisfies **S** then (I, ι) , with the topology defined in §4, is an M -coalgebra in **Top**, and indeed the universal solution in **Top**.

The proof involves an analysis of equality in Ia , that is, of the possible representations in $\mathcal{I}a$ of a given element of Ia .

Lemma 6.1 *For each $a \in \mathbb{A}$ and $n \in \mathbb{N}$, the category $\mathcal{I}_n a$ is finite.*

Proof Follows from finiteness of M . $\square$

The last paragraph of §4 shows that there may be objects τ, τ' of $\mathcal{I}a$ such that $\text{pr}_n(\tau)$ and $\text{pr}_n(\tau')$ are in the same connected-component of $\mathcal{I}_n a$ for all n but τ and τ' are in different connected-components of $\mathcal{I}a$. However, τ and τ' are in the same component if $\text{pr}_n(\tau)$ and $\text{pr}_n(\tau')$ can be connected by a diagram of the same shape for each n :

Lemma 6.2 (Equality in Ia) *Let $a \in \mathbb{A}$ and $\tau, \tau' \in \mathcal{I}a$. Then $[\tau] = [\tau'] \in Ia$ if and only if there exist a finite connected category $\mathbb{K}$, objects $k, k' \in \mathbb{K}$, and for each $n \geq 1$, a functor $F_n : \mathbb{K} \longrightarrow \mathcal{I}_n a$ such that $F_n k = \text{pr}_n \tau$ and $F_n k' = \text{pr}_n \tau'$.*

Proof ‘Only if’ is simple. For ‘if’ we use Lemma 5.2. Take $\mathbb{J}$ to be the discrete category on two objects j, j' , define $P : \mathbb{J} \longrightarrow \mathbb{K}$ by $P(j) = k$ and $P(j') = k'$, take $\mathcal{L}$ to be the limit $\mathcal{I}a$ of the categories $\mathcal{L}_n = \mathcal{I}_n a$, and take $D(j) = \tau$ and $D(j') = \tau'$. Then for each $n \geq 1$, the set of factorizations in Lemma 5.2 is nonempty by hypothesis, and finite since $\mathbb{K}$ and $\mathcal{I}_n a$ are finite, so there is a functor $G : \mathbb{K} \longrightarrow \mathcal{I}a$ such that $Gk = \tau$ and $Gk' = \tau'$. But $\mathbb{K}$ is connected, so $[\tau] = [\tau']$. $\square$

Assume from now on that $(\mathbb{A}, M)$ satisfies **S**.

Each module element $q : d \multimap c$ induces a function $\phi_q : Id \longrightarrow Ic$ by

$$\phi_q[\cdots \xrightarrow{r_2} d_1 \xrightarrow{r_1} d] = [\cdots \xrightarrow{r_2} d_1 \xrightarrow{r_1} d \xrightarrow{q} c],$$

or equivalently,

$$\phi_q = \left(Id \xrightarrow{q \otimes -} (M \otimes I)c \xrightarrow[\sim]{\iota_c^{-1}} Ic \right)$$

where the first map is the coprojection

$$\text{copr}_q = q \otimes - : Id \longrightarrow \int^{d'} M(d', c) \times Id' = (M \otimes I)c.$$

For each $a \in \mathbb{A}$, the topology on Ia is generated by taking $\phi_{p_1} \phi_{p_2} \cdots \phi_{p_n}(Ib_n)$ to be a closed subset of Ia whenever $n \in \mathbb{N}$ and

$$(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_2} b_1 \xrightarrow{p_1} b_0) \in \mathcal{I}_n a.$$

It is shown in the following pages that (I, ι) is an M -coalgebra in **Top**, and along the way that each Ia is compact Hausdorff.

We start with the Hausdorff property. Define, for each $n \in \mathbb{N}$ and $a \in \mathbb{A}$, a binary relation R_n^a on Ia by

$$R_n^a = \bigcup \{ (\phi_{p_1} \cdots \phi_{p_n}(Ib_n))^2 \mid (b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \in \mathcal{I}_n a \} \subseteq Ia \times Ia.$$

Equivalently, $(t, t') \in R_n^a$ when there exists $(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0)$ such that t and t' can both be written in the form

$$[\cdots \xrightarrow{\quad} \cdot \xrightarrow{\quad} b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0].$$

As a subset of $Ia \times Ia$, R_n^a is closed (by finiteness of $\mathcal{I}_n a$). As a relation, R_n^a is reflexive and symmetric, but not in general transitive: for instance, in the Freyd self-similarity system, $R_1^1 = [0, \frac{1}{2}]^2 \cup [\frac{1}{2}, 1]^2 \subseteq [0, 1]^2$.

For any set S , write $\Delta_S = \{(s, s) \mid s \in S\} \subseteq S \times S$.

Lemma 6.3 (Relations determine equality) $\bigcap_{n \in \mathbb{N}} R_n^a = \Delta_{Ia}$ for each a .

Proof Certainly $\bigcap_{n \in \mathbb{N}} R_n^a \supseteq \Delta_{Ia}$. Conversely, let $(t, t') \in \bigcap_{n \in \mathbb{N}} R_n^a$, writing

$$\begin{aligned} t &= [\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0 = a], \\ t' &= [\cdots \xrightarrow{m'_2} a'_1 \xrightarrow{m'_1} a'_0 = a]. \end{aligned}$$

For each $n \in \mathbb{N}$, we may choose $(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_1) \in \mathcal{I}_n a$ such that $t, t' \in \phi_{p_1} \cdots \phi_{p_n}(Ib_n)$. Since $\mathcal{I}a$ is componentwise cofiltered, there is for each $n \in \mathbb{N}$ a span in $\mathcal{I}a$ of the form

$$\begin{array}{ccccccc} \cdots & \xrightarrow{m_{n+2}} & a_{n+1} & \xrightarrow{m_{n+1}} & a_n & \xrightarrow{m_n} & \cdots & \xrightarrow{m_1} & a_0 = a \\ & & \uparrow & & \uparrow & & & & \uparrow 1_a \\ \cdots & \xrightarrow{\quad} & \cdot & \xrightarrow{\quad} & \cdot & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & a \\ & & \downarrow & & \downarrow & & & & \downarrow 1_a \\ \cdots & \xrightarrow{\quad} & \cdot & \xrightarrow{\quad} & b_n & \xrightarrow{p_n} & \cdots & \xrightarrow{p_1} & b_0 = a, \end{array}$$

hence a span in $\mathcal{I}_n a$ of the form

$$\begin{array}{ccc} & \cdot & \\ \swarrow & & \searrow \\ (a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0) & & (b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0). \end{array}$$

The same is true for t' , so for each $n \in \mathbb{N}$ there is a diagram of the form

$$\begin{array}{ccccc} & & \cdot & & \cdot \\ \swarrow & & \searrow & & \swarrow \searrow \\ (a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0) & & \cdot & & (a'_n \xrightarrow{m'_n} \cdots \xrightarrow{m'_1} a'_0) \end{array}$$

in $\mathcal{I}_n a$. So by Lemma 6.2, taking $\mathbb{K}$ to be the evident category with four non-identity arrows, $t = t'$. $\square$

Proposition 6.4 (*Ia Hausdorff*) Ia is Hausdorff for all $a \in \mathbb{A}$.

Proof Δ_{Ia} is the intersection of the closed subsets R_n^a of $Ia \times Ia$. $\square$

The next step is to consider $I : \mathbb{A} \longrightarrow \mathbf{Set}$ as a quotient of $\text{ob } \mathcal{I}$, the composite of $\mathcal{I} : \mathbb{A} \longrightarrow \mathbf{Cat}$ with the objects functor $\text{ob} : \mathbf{Cat} \longrightarrow \mathbf{Set}$. This functor $\text{ob } \mathcal{I}$ is nondegenerate (even if **S** does not hold), since

$$\text{ob } \mathcal{I} \cong \sum_b \text{ob}(\mathcal{I}b) \times M(b, -)$$

and the class of nondegenerate functors is closed under sums. Moreover, $\text{ob } \mathcal{I}$ carries an M -coalgebra structure ι , given by the usual formula

$$\iota_{a_0}(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0) = m_1 \otimes (\cdots \xrightarrow{m_2} a_1)$$

or equivalently by taking ι_a to be the composite

$$\text{ob}(\mathcal{J}a) \xrightarrow{\sim} \sum_b M(b, a) \times \text{ob}(\mathcal{J}b) \xrightarrow{\text{canonical}} \int^b M(b, a) \times \text{ob}(\mathcal{J}b). \quad (30)$$

So $(\text{ob} \mathcal{J}, \iota)$ is a coalgebra in **Set**, and there is a canonical map of coalgebras $\pi : (\text{ob} \mathcal{J}, \iota) \longrightarrow (I, \iota)$.

This construction can be topologized. For each $a \in \mathbb{A}$, the set $\text{ob}(\mathcal{J}a)$ is the limit of the diagram

$$\cdots \longrightarrow \text{ob}(\mathcal{J}_2 a) \longrightarrow \text{ob}(\mathcal{J}_1 a),$$

and giving each $\text{ob}(\mathcal{J}_n a)$ the discrete topology induces a topology on $\text{ob}(\mathcal{J}a)$; in this way, $\text{ob} \mathcal{J}$ becomes a functor $\mathbb{A} \longrightarrow \mathbf{Top}$. Each $\mathcal{J}_n a$ is finite, so each $\text{ob}(\mathcal{J}a)$ is compact Hausdorff, so $\text{ob} \mathcal{J} : \mathbb{A} \longrightarrow \mathbf{Top}$ is nondegenerate. Moreover, the maps ι_a are continuous, since in (30) the first map is a homeomorphism and the second is a quotient map. So $(\text{ob} \mathcal{J}, \iota)$ is a coalgebra in **Top**. We will see that the maps $\pi_a : \text{ob}(\mathcal{J}a) \longrightarrow Ia$ exhibit each Ia as not merely a set-theoretic quotient of $\text{ob}(\mathcal{J}a)$, but a topological quotient.

For each $a \in \mathbb{A}$, the space $\text{ob}(\mathcal{J}a)$ is compact. It is also **second countable** (has a countable basis of open sets). Hence:

Lemma 6.5 *Every sequence in $\text{ob}(\mathcal{J}a)$ has a convergent subsequence.* □

The following result must hold if the spaces Ia are to be compact.

Lemma 6.6 (Intersections of basic closed sets) *Let $a \in \mathbb{A}$, $n \in \mathbb{N}$,*

$$(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0) \in \mathcal{J}a, \quad (b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \in \mathcal{J}_n a.$$

Then

$$[\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0] \in \phi_{p_1} \cdots \phi_{p_n}(Ib_n)$$

if and only if

$$\forall r \in \mathbb{N}, \phi_{m_1} \cdots \phi_{m_r}(Ia_r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \neq \emptyset.$$

Proof ‘Only if’ is trivial. For ‘if’, write $\alpha = (\cdots \xrightarrow{m_1} a_0)$. By hypothesis, we may choose for each $r \in \mathbb{N}$ objects

$$\begin{aligned} \alpha_r &= (\cdots \xrightarrow{m_{r+2}^r} a_{r+1}^r \xrightarrow{m_{r+1}^r} a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0), \\ \beta_r &= (\cdots \xrightarrow{p_{n+2}^r} b_{n+1}^r \xrightarrow{p_{n+1}^r} b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \end{aligned}$$

of $\mathcal{J}a$ such that $[\alpha_r] = [\beta_r]$. By Proposition 5.5, there is for each $r \in \mathbb{N}$ a span

$$\alpha_r \longleftarrow \cdot \longrightarrow \beta_r$$

in $\mathcal{I}a$. By Lemma 6.5, $(\beta_r)_{r \in \mathbb{N}}$ has a subsequence $(\beta_{r_k})_{k \in \mathbb{N}}$ convergent to β , say, and β is of the form

$$\cdots \xrightarrow{p_{n+1}} b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0.$$

For each $r \in \mathbb{N}$, we may choose $k \in \mathbb{N}$ such that $r_k \geq r$ and $\text{pr}_r(\beta_{r_k}) = \text{pr}_r(\beta)$; we then have a span in $\mathcal{I}a$ of the form

$$\alpha_{r_k} \longleftarrow \cdot \longrightarrow \beta_{r_k},$$

hence, applying pr_r , a span in $\mathcal{I}_r a$ of the form

$$\text{pr}_r(\alpha) \longleftarrow \cdot \longrightarrow \text{pr}_r(\beta).$$

So by Lemma 6.2, $[\alpha] = [\beta] \in \phi_{p_1} \cdots \phi_{p_n}(Ib_n)$. $\square$

Proposition 6.7 (*Ia as a quotient*) *For each $a \in \mathbb{A}$, the canonical surjection $\pi_a : \text{ob}(\mathcal{I}a) \longrightarrow Ia$ is a topological quotient map.*

Proof First, π_a is continuous. Let $n \in \mathbb{N}$ and $(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \in \mathcal{I}_n a$; we must show that $\pi_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n))$ is a closed subset of $\text{ob}(\mathcal{I}a)$. By Lemma 6.6, an element $(\cdots \xrightarrow{m_1} a_0)$ of $\text{ob}(\mathcal{I}a)$ belongs to this subset if and only if

$$\phi_{m_1} \cdots \phi_{m_r}(Ia_r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \neq \emptyset \quad (31)$$

for all $r \in \mathbb{N}$. In other words,

$$\pi_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n)) = \bigcap_{r \in \mathbb{N}} \text{pr}_r^{-1} W_r$$

where $\text{pr}_r : \mathcal{I}a \longrightarrow \mathcal{I}_r a$ and $W_r \subseteq \text{ob}(\mathcal{I}_r a)$ is the set of elements $(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0)$ satisfying (31). But each $\text{ob}(\mathcal{I}_r a)$ is discrete and each pr_r continuous, so $\bigcap_{r \in \mathbb{N}} \text{pr}_r^{-1} W_r$ is closed, as required.

Second, π_a is closed, since $\text{ob}(\mathcal{I}a)$ is compact and Ia is Hausdorff; and any continuous closed surjection is a quotient map. $\square$

Corollary 6.8 (*Ia compact*) *Ia is compact for all $a \in \mathbb{A}$.* $\square$

Corollary 6.9 (*Properties of If*) *For each map $a \xrightarrow{f} a'$ in $\mathbb{A}$, the map $Ia \xrightarrow{If} Ia'$ is continuous and closed.*

Proof There is a commutative square

$$\begin{array}{ccc} \text{ob}(\mathcal{I}a) & \xrightarrow{\pi_a} & Ia \\ \text{ob}(\mathcal{I}f) \downarrow & & \downarrow If \\ \text{ob}(\mathcal{I}a') & \xrightarrow{\pi_{a'}} & Ia' \end{array}$$

in which π_a is a topological quotient map and $\text{ob}(\mathcal{I}f)$ and $\pi_{a'}$ are continuous, so If is also continuous. Ia is compact and Ia' Hausdorff, so If is closed. $\square$

Corollary 6.10 (ι_a continuous) *For each $a \in \mathbb{A}$, the map $\iota_a : Ia \longrightarrow (M \otimes I)a$ is continuous.*

Proof As for the previous lemma, using the square

$$\begin{array}{ccc} \text{ob}(\mathcal{J}a) & \xrightarrow{\pi_a} & Ia \\ \downarrow \iota_a & & \downarrow \iota_a \\ (M \otimes \text{ob } \mathcal{J})a & \xrightarrow{(M \otimes \pi)_a} & (M \otimes I)a. \end{array}$$

□

Corollary 6.11 (I, ι) is an M -coalgebra in **Top**.

□

Our final task is to prove that for any M -coalgebra (X, ξ) in **Top**, the unique map $\bar{\xi} : (X, \xi) \longrightarrow (I, \iota)$ of coalgebras in **Set** is continuous. To do this we show that the inverse image $\bar{\xi}_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n))$ of each basic closed set is closed. This inverse image is larger than it might appear. For, given $x \in Xa$, one might imagine that if $\bar{\xi}_a(x) \in \phi_{p_1} \cdots \phi_{p_n}(Ib_n)$ then x can be resolved along some diagram of the form

$$\cdots \dashrightarrow \cdot \dashrightarrow b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0;$$

but while this is certainly a sufficient condition, it is not necessary:

Example 6.12 Let $(\mathbb{A}, M)$ be the Freyd self-similarity system. Choose an endpoint-preserving continuous map $\xi_1 : [0, 1] \longrightarrow [0, 2]$ satisfying $\xi_1(2/3) = 2/3$; this gives an M -coalgebra structure ξ on $X = (\{\star\} \xrightleftharpoons[1]{0} [0, 1])$. Now $2/3 \in X1$ has a resolution along

$$\cdots \dashrightarrow [0, \tfrac{1}{2}] \dashrightarrow 1 \dashrightarrow [0, \tfrac{1}{2}] \dashrightarrow 1,$$

so $\bar{\xi}_1(2/3) = 0$, so $\bar{\xi}_1(2/3) \in \phi_{m_1}(I0)$ where $m_1 = 0 : 0 \dashrightarrow 1$. But $2/3$ has no resolution ending in m_1 .

To describe this inverse image we need some notation. Given an M -coalgebra $X = (X, \xi)$ in **Top**, $r \in \mathbb{N}$, and $(a_r \dashrightarrow \cdots \dashrightarrow a_0) \in \mathcal{J}_r a$, let

$$\text{Res}_X(a_r \dashrightarrow \cdots \dashrightarrow a_0) \subseteq Xa$$

be the set of all $x \in Xa$ having a resolution along some diagram of the form

$$\cdots \dashrightarrow \cdot \dashrightarrow a_r \dashrightarrow \cdots \dashrightarrow a_0.$$

Lemma 6.13 (Inverse image) *Let $a \in \mathbb{A}$, $n \in \mathbb{N}$, and $(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \in \mathcal{I}_n a$; let (X, ξ) be an M -coalgebra in **Set** and $x \in Xa$. Then $x \in \bar{\xi}_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n))$ if and only if*

for all $r \in \mathbb{N}$, there exists $(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0) \in \mathcal{I}_r a$ such that $x \in \text{Res}_X(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0)$ and $\phi_{m_1} \cdots \phi_{m_r}(Ia_r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \neq \emptyset$.

Proof For ‘only if’, choose $(\cdots \xrightarrow{m_1} a_0) \in \mathcal{I}a$ along which x can be resolved. Then

$$[\cdots \xrightarrow{m_1} a_0] = \bar{\xi}_a(x) \in \phi_{p_1} \cdots \phi_{p_n}(Ib_n)$$

and the result follows.

For ‘if’, choose for each $r \in \mathbb{N}$ an element $(\cdots \xrightarrow{m_1^r} a_0^r) \in \mathcal{I}a$ along which x can be resolved and such that

$$\phi_{m_1^r} \cdots \phi_{m_r^r}(Ia_r^r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \neq \emptyset.$$

By compactness of Ia ,

$$\bigcap_{r \in \mathbb{N}} \phi_{m_1^r} \cdots \phi_{m_r^r}(Ia_r^r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \quad (32)$$

is nonempty. But $\bigcap_{r \in \mathbb{N}} \phi_{m_1^r} \cdots \phi_{m_r^r}(Ia_r^r)$ has cardinality at most 1 by Lemma 6.3, and contains $\bar{\xi}_a(x)$ since x can be resolved along $(\cdots \xrightarrow{m_1^r} a_0^r)$ for each r , so the set (32) is $\{\bar{\xi}_a(x)\}$. The result follows. $\square$

This says what the inverse images of the basic closed sets are, and we now prepare to show that they are closed.

Lemma 6.14 (Coprojections closed) *Let $X : \mathbb{A} \longrightarrow \mathbf{Top}$ be a nondegenerate functor and $m : b \dashrightarrow a$ a module element. Then the coprojection*

$$m \otimes - : Xb \longrightarrow (M \otimes X)a$$

is closed.

Proof Trivially, $m \otimes -$ is the composite

$$Xb \xrightarrow{(m, -)} \sum_{b'} M(b', a) \times Xb' \xrightarrow{\text{canonical}} \int^{b'} M(b', a) \times Xb' = (M \otimes X)a.$$

The first map is closed since it is a coproduct-coprojection, and the second is closed by Lemma 3.3. $\square$

Lemma 6.15 *Let (X, ξ) be an M -coalgebra in **Top**, $r \in \mathbb{N}$, and $(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0) \in \mathcal{I}_r a$. Then $\text{Res}_X(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0)$ is a closed subset of Xa .*

Proof When $r = 0$ this is trivial. Suppose inductively that the result holds for $r \in \mathbb{N}$, and let $(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_1} a_0) \in \mathcal{J}_{r+1}a$. An element $x \in Xa$ is in $\text{Res}_X(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_1} a_0)$ if and only if there exists $x_1 \in Xa_1$ such that

$$\xi_a(x) = m_1 \otimes x_1 \quad \text{and} \quad x_1 \in \text{Res}_X(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_2} a_1),$$

so

$$\text{Res}_X(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_1} a_0) = \xi_a^{-1} \left(m_1 \otimes \text{Res}_X(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_2} a_1) \right).$$

But $\text{Res}_X(a_{r+1} \xrightarrow{m_{r+1}} \cdots \xrightarrow{m_2} a_1)$ is closed by inductive hypothesis, $m_1 \otimes -$ is closed by Lemma 6.14, and ξ_a is continuous, so the induction is complete. $\square$

Theorem 6.16 (Universal solution in $\mathbf{Top}$) *(I, ι) is the universal solution of $(\mathbb{A}, M)$ in $\mathbf{Top}$.*

Proof It remains to show that for any M -coalgebra (X, ξ) in $\mathbf{Top}$, the unique map $\bar{\xi} : (X, \xi) \longrightarrow (I, \iota)$ is continuous in each component. So let $a \in \mathbb{A}$, $n \in \mathbb{N}$, and $(b_n \xrightarrow{p_n} \cdots \xrightarrow{p_1} b_0) \in \mathcal{J}_n a$; we must show that

$$\bar{\xi}_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n)) \subseteq Xa$$

is closed. For each $r \in \mathbb{N}$, write

$$V_r = \bigcup \text{Res}_X(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0)$$

where the union is over all $(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0) \in \mathcal{J}_r a$ such that

$$\phi_{m_1} \cdots \phi_{m_r}(Ia_r) \cap \phi_{p_1} \cdots \phi_{p_n}(Ib_n) \neq \emptyset.$$

Then Lemma 6.13 says that $\bar{\xi}_a^{-1}(\phi_{p_1} \cdots \phi_{p_n}(Ib_n)) = \bigcap_{r \in \mathbb{N}} V_r$, so it is enough to prove that each V_r is closed, and this follows from Lemmas 6.1 and 6.15. $\square$

A Appendix: Solvability

Here we finish the proof of

Theorem A.1 (Existence of universal solution) *The following are equivalent for a self-similarity system $(\mathbb{A}, M)$:*

- a. $(\mathbb{A}, M)$ satisfies **S**
- b. $(\mathbb{A}, M)$ has a universal solution in **Top**

c. $(\mathbb{A}, M)$ has a universal solution in **Set**.

We proved (a) $\Rightarrow$ (b) in §6 and (b) $\Rightarrow$ (c) as Proposition 3.6, so it remains to prove (c) $\Rightarrow$ (a).

Fix a self-similarity system $(\mathbb{A}, M)$. In this appendix, ‘ M -coalgebra’ means ‘ M -coalgebra in **Set**’.

We begin by constructing representable-type coalgebras and proving a Yoneda-type lemma. Take $(a_\bullet, m_\bullet) = (\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0) \in \mathbb{E}(\mathcal{J})$. Then there is a functor

$$H^{(a_\bullet, m_\bullet)} = \sum_{n \in \mathbb{N}} \mathbb{A}(a_n, -) : \mathbb{A} \longrightarrow \mathbf{Set},$$

and any representable functor is flat, so $H^{(a_\bullet, m_\bullet)}$ is nondegenerate by Theorem 2.6. Also

$$(M \otimes H^{(a_\bullet, m_\bullet)})b \cong \sum_{n \in \mathbb{N}} (M \otimes \mathbb{A}(a_n, -))b \cong \sum_{n \in \mathbb{N}} M(a_n, b),$$

so an M -coalgebra structure on $H^{(a_\bullet, m_\bullet)}$ amounts to a natural transformation

$$\sum_{n \in \mathbb{N}} \mathbb{A}(a_n, -) \longrightarrow \sum_{n \in \mathbb{N}} M(a_n, -).$$

There is a unique such transformation sending 1_{a_n} to $m_{n+1} \in M(a_{n+1}, a_n)$ for each $n \in \mathbb{N}$; let $\theta^{(a_\bullet, m_\bullet)}$ be the corresponding coalgebra structure on $H^{(a_\bullet, m_\bullet)}$.

This defines an M -coalgebra $(H^{(a_\bullet, m_\bullet)}, \theta^{(a_\bullet, m_\bullet)})$ for each object $(a_\bullet, m_\bullet)$ of $\mathbb{E}(\mathcal{J})$. In fact there is a functor

$$(H^\bullet, \theta^\bullet) : \mathbb{E}(\mathcal{J})^{\text{op}} \longrightarrow \mathbf{Coalg}(M, \mathbf{Set}),$$

since any map $f_\bullet : (a_\bullet, m_\bullet) \longrightarrow (a'_\bullet, m'_\bullet)$ in $\mathbb{E}(\mathcal{J})$ induces a map

$$H^{(a'_\bullet, m'_\bullet)} = \sum_{n \in \mathbb{N}} \mathbb{A}(a'_n, -) \xrightarrow{\sum f_n^*} \sum_{n \in \mathbb{N}} \mathbb{A}(a_n, -) = H^{(a_\bullet, m_\bullet)}$$

respecting the coalgebra structures.

Lemma A.2 (‘Yoneda’) *There is a bijection*

$$\begin{aligned} & \mathbf{Coalg}(M, \mathbf{Set}) \left((H^{(a_\bullet, m_\bullet)}, \theta^{(a_\bullet, m_\bullet)}), (X, \xi) \right) \\ & \cong \{ \text{resolutions along } (a_\bullet, m_\bullet) \text{ in } (X, \xi) \} \end{aligned} \quad (33)$$

natural in $(a_\bullet, m_\bullet) \in \mathbb{E}(\mathcal{J})$ and $(X, \xi) \in \mathbf{Coalg}(M, \mathbf{Set})$. More precisely, if $x \in Xa_0$ then the coalgebra maps $H^{(a_\bullet, m_\bullet)} \longrightarrow X$ sending 1_{a_0} to x correspond to the resolutions of x along $(a_\bullet, m_\bullet)$.

Proof A natural transformation $\alpha : H^{(a_\bullet, m_\bullet)} \longrightarrow X$ amounts to a sequence $(x_n)_{n \in \mathbb{N}}$ with $x_n \in Xa_n$, by the standard Yoneda Lemma. It is a map of coalgebras if and only if

$$\begin{array}{ccc} \sum_{n \in \mathbb{N}} \mathbb{A}(a_n, -) & \xrightarrow{\alpha} & X \\ \theta^{(a_\bullet, m_\bullet)} \downarrow & & \downarrow \xi \\ \sum_{n \in \mathbb{N}} M(a_n, -) & \xrightarrow{M \otimes \alpha} & M \otimes X \end{array}$$

commutes, if and only if this diagram commutes when we take 1_{a_n} at the top-left corner for every $n \in \mathbb{N}$. We have

$$\begin{array}{ccc} 1_{a_n} & \xrightarrow{\quad} & x_n \\ \downarrow & & \downarrow \xi(x_n) \\ m_{n+1} & \xrightarrow{\quad} & m_{n+1} \otimes x_{n+1}, \end{array}$$

so α is a map of coalgebras just when $\xi(x_n) = m_{n+1} \otimes x_{n+1}$ for all $n \in \mathbb{N}$. A coalgebra map $(H^{(a_\bullet, m_\bullet)}, \theta^{(a_\bullet, m_\bullet)}) \longrightarrow (X, \xi)$ therefore amounts to a sequence $(x_n)_{n \in \mathbb{N}}$ satisfying $\xi(x_n) = m_{n+1} \otimes x_{n+1}$ for all n , that is, a resolution along $(a_\bullet, m_\bullet)$ in (X, ξ) . This establishes the bijection (33); naturality follows from the naturality in the standard Yoneda Lemma. $\square$

We have met one other M -coalgebra: $(\text{ob } \mathcal{J}, \iota)$, constructed after Proposition 6.4. (Recall from §5 that (I, ι) is only known to be a (nondegenerate) coalgebra if **S** holds.)

Corollary A.3 (Tautological map) *For each $(a_\bullet, m_\bullet) \in \mathbb{E}(\mathcal{J})$ there is a canonical map of M -coalgebras*

$$\kappa^{(a_\bullet, m_\bullet)} : (H^{(a_\bullet, m_\bullet)}, \theta^{(a_\bullet, m_\bullet)}) \longrightarrow (\text{ob } \mathcal{J}, \iota)$$

sending 1_{a_0} to $(a_\bullet, m_\bullet)$.

Proof Every $(a_\bullet, m_\bullet) \in \mathbb{E}(\mathcal{J})$, regarded as an element of $(\text{ob } \mathcal{J})_{a_0}$, has a tautological resolution in $\text{ob } \mathcal{J}$. By Lemma A.2, the corresponding map $\kappa^{(a_\bullet, m_\bullet)}$ of coalgebras sends 1_{a_0} to $(a_\bullet, m_\bullet)$. $\square$

We can now finish the proof of Theorem A.1.

Suppose that $(\mathbb{A}, M)$ has a universal solution (J, γ) . Then there is a map $\beta : (\text{ob } \mathcal{J}, \iota) \longrightarrow (J, \gamma)$ of M -coalgebras. I claim that the natural transformation β can be factorized as

$$\begin{array}{ccc} \text{ob } \mathcal{J} & \xrightarrow{\beta} & J, \\ \pi \searrow & & \nearrow \bar{\beta} \\ & I & \end{array}$$

or equivalently that for each $a \in \mathbb{A}$, the function $\beta_a : \text{ob}(\mathcal{J}a) \longrightarrow \mathcal{J}a$ is constant on connected-components of $\mathcal{J}a$, or equivalently that if $f_\bullet : (a_\bullet, m_\bullet) \longrightarrow (b_\bullet, p_\bullet)$ in $\mathcal{J}a$ then $\beta_a(a_\bullet, m_\bullet) = \beta_a(b_\bullet, p_\bullet)$. Indeed, given such an $f_\bullet$, there are coalgebra maps

$$\begin{array}{ccc} (H^{(a_\bullet, m_\bullet)}, \theta^{(a_\bullet, m_\bullet)}) & \xrightarrow{\kappa^{(a_\bullet, m_\bullet)}} & (\text{ob } \mathcal{J}, \iota) \\ f_\bullet^* \uparrow & & \searrow \kappa^{(b_\bullet, p_\bullet)} \\ (H^{(b_\bullet, p_\bullet)}, \theta^{(b_\bullet, p_\bullet)}) & \xrightarrow{\kappa^{(b_\bullet, p_\bullet)}} & (\text{ob } \mathcal{J}, \iota) \end{array} \xrightarrow{\beta} (J, \gamma)$$

and $\beta \circ \kappa^{(a_\bullet, m_\bullet)} \circ f_\bullet^* = \beta \circ \kappa^{(b_\bullet, p_\bullet)}$ by terminality of (J, γ) . But

$$\begin{aligned} \kappa^{(a_\bullet, m_\bullet)} f_\bullet^*(1_a) &= \kappa^{(a_\bullet, m_\bullet)}(1_a \circ f_0) = \kappa^{(a_\bullet, m_\bullet)}(1_a) = (a_\bullet, m_\bullet), \\ \kappa^{(b_\bullet, p_\bullet)}(1_a) &= (b_\bullet, p_\bullet), \end{aligned}$$

so $\beta_a(a_\bullet, m_\bullet) = \beta_a(b_\bullet, p_\bullet)$ as required.

The transformation $\bar{\beta} : I \longrightarrow J$ induces a functor $\mathbb{E}(I) \longrightarrow \mathbb{E}(J)$ over $\mathbb{A}$. Moreover, there is a natural transformation $\text{id}_{\mathbf{Cat}} \longrightarrow D\Pi_0$ (the unit of the adjunction $\Pi_0 \dashv D$), hence a transformation $\mathcal{J} \longrightarrow D\Pi_0\mathcal{J} = DI$, hence a functor $\mathbb{E}(\mathcal{J}) \longrightarrow \mathbb{E}(DI) \cong \mathbb{E}(I)$ over $\mathbb{A}$. So there is a commutative triangle

$$\begin{array}{ccc} \mathbb{E}(\mathcal{J}) & \xrightarrow{F} & \mathbb{E}(J) \\ \text{pr} \searrow & & \swarrow \text{pr} \\ & \mathbb{A} & \end{array}$$

in $\mathbf{Cat}$, where pr denotes a projection. Now, condition **S** says that if $\mathbb{K}$ is either of the categories $(\bullet \longrightarrow \bullet \longleftarrow \bullet)$ or $(\bullet \rightrightarrows \bullet)$ and if $G : \mathbb{K} \longrightarrow \mathbb{E}(\mathcal{J})$ then $\text{pr} \circ G$ admits a cone. But given such a G , nondegeneracy of J implies that $F \circ G$ admits a cone, so $\text{pr} \circ G = \text{pr} \circ F \circ G$ admits a cone, as required.

References

- [ABLR] Jiří Adámek, Francis Borceux, Stephen Lack, Jiří Rosický, A classification of accessible categories, *Journal of Pure and Applied Algebra* 175 (2002), 7–30.
- [Age] P. Ageron, Catégories accessibles à produits fibrés, *Diagrammes* 36 (1996), 1–11.
- [Barr] Michael Barr, Terminal coalgebras in well-founded set theory, *Theoretical Computer Science* 114 (1993), no. 2, 299–315.
- [Bor] Francis Borceux, *Handbook of Categorical Algebra 1: Basic Category Theory*, Encyclopedia of Mathematics and its Applications 50, Cambridge University Press, 1994.
- [Bou] Nicolas Bourbaki, *General Topology* Part 1, Springer, 1966.
- [Fal] Kenneth Falconer, *Fractal Geometry*, John Wiley & Sons, 1990.
- [Fre] Peter Freyd, Real coalgebra, post on categories mailing list, 22 December 1999, available via www.mta.ca/~cat-dist

- [GU] P. Gabriel, F. Ulmer, Lokal präsentierbare Kategorien, Lecture Notes in Mathematics 221, Springer, 1971.
- [GV] A. Grothendieck, J.L. Verdier, Préfaisceaux, in Artin, Grothendieck, Verdier (eds.), *Théorie des topos et cohomologie étale des schémas (SGA4)*, Lecture Notes in Mathematics 269, Springer, 1972.
- [Hut] John E. Hutchinson, Fractals and self-similarity, *Indiana University Mathematics Journal* 30 (1981), no. 5, 713–747.
- [Joy] A. Joyal, Disks, duality and Θ -categories, preprint, 1997.
- [Kön] Dénes König, Sur les correspondences multivoques des ensembles, *Fundamenta Mathematicae* 8 (1926), 114–134.
- [Lai] C. Lair, Sur les genres d’esquissabilité des catégories modelables (accessibles) possédant les limites d’indexation finies [resp. finies et non vides, finies et connexes, finies et connexes et non vides], *Diagrammes* 35 (1996), 53–90.
- [Lam] Joachim Lambek, A fixpoint theorem for complete categories, *Mathematische Zeitschrift* 103 (1968), 151–161.
- [Lei1] Tom Leinster, A survey of definitions of n -category, e-print math.CT/0107188, 2001; also *Theory and Applications of Categories* 10 (2002), no. 1, 1–70.
- [Lei2] Tom Leinster, A general theory of self-similarity II: recognition, e-print math.DS/0411345, 2004.
- [Lei3] Tom Leinster, General self-similarity: an overview, e-print math.DS/0411343, 2004.
- [LM] Douglas Lind, Brian Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, 1995.
- [Mac] Saunders Mac Lane, *Categories for the Working Mathematician*, Graduate Texts in Mathematics 5, Springer, 1971.
- [MM] Saunders Mac Lane, Ieke Moerdijk, *Sheaves in Geometry and Logic*, Springer, 1994.
- [Mil] John Milnor, *Dynamics in One Complex Variable*, Vieweg, 1999.
- [PR] H.-O. Peitgen, P.H. Richter, *The Beauty of Fractals*, Springer, 1986.
- [Web] Helmut Weberpals, Vertauschbarkeit von Limites und Colimites, *Manuscripta Mathematica* 49 (1984), no. 1, 9–26.